

# PRODUCT CATALOGUE

2023.



***Belupo is the second-largest producer of prescription drugs, OTC products and enteral food in Croatia.***



According to the anatomical-therapeutic-chemical (ATC) classification, Belupo is present in 12 ATK groups- including enteral nutrition.

Belupo's product portfolio is focused on the following:

cardiovascular system medicines, musculo-skeletal system medicines, nervous system medicines, gastrointestinal system medicines, dermatological and general anti-infective medicines for systemic use.

Development and production of high quality products is the main strategic orientation for all 1300 Belupo's employees, both on domestic and 18 European markets.

After many years of constant growth and development, in 2017. Belupo has finished investment cycle - building a new solid, semi-liquid and liquid medicine factory. These are Belupo's factories for the future, the future which is created by knowledge of their employees and by implementing new technologies in the manufacture of generic drugs with added value, and new generation medicines as well.



New factories were the biggest greenfield investment in Croatia, in which was invested more than 70 million Euros. With the modern technology and processes organization according to latest standards of good consumer practice, these factories represent one of the most modern pharmaceutical plants in the region. New production lines increase production capacities by 150 percent and ensure annual production of 100 million packaging.

The new plants meet top standards of modern pharmaceutical industry, including the Good Manufacturing Practice Standards, The European Pharmaceutical Inspection Convention regulations, and the requirements and standards set in the Croatian Drug Law.

Numerous and diverse forms of cooperation include reputable foreign business partners: Abbott, Pharmathen, Medis, Lupin and Torrent confirm and guarantee our dedication to latest developments in the field of field of pharmaceuticals.

# Products

*The wide range of Belupo's products is categorized under three main groups: prescription medicines, OTC products and food for special medical purposes.*



Every year we are upgrading our portfolio by developing and manufacturing new products medicinal products, food supplements, cosmetic products and medical devices, as well as by improving the existing ones.

We are proud to say that nowadays our product portfolio includes more than 1660 products from 164 active ingredients, packaged in 345 different dose sizes and 189 pharmaceutical dosage forms. Most of them are a result of our knowledge in research and development, production and quality control.

We are producing generic medicinal products that are not inferior to most leading pharmaceutical brands in terms of quality, safety and efficacy.



# Broad range of therapeutic areas

## Prescription medicines



- alimentary tract and metabolism (group A),
- blood and blood forming organs (group B),
- cardiovascular system (group C),
- dermatologicals (group D),
- genito-urinary system and sex hormones (group G),
- antiinfectives for systemic use (group J),
- antineoplastic and immunomodulating agents (group L),
- musculo-skeletal system (group M),
- nervous system (group N),
- antiparasitic products, insecticides and repellents (group P),
- respiratory system (group R).

### **Food for special medical purposes**

- NUTRIXA - taste: vanilla, chocolate, coffee, strawberry, peach
- NUTRIBEL COMPLEX - taste: chocolate, cappuccino, orange-biscuit
- NUTRIBEL DIABET - taste: chocolate, vanilla, strawberry

## OTC



- herbal medicines
- food supplements
- cosmetics
- non – prescription medicines
- medical Device
- auxiliary substances

# Pharmaceuticals

## A

### ALIMENTARY TRACT AND METABOLISM

#### AGNIS

##### VILDAGLIPTIN

Vildagliptin is an oral anti-hyperglycemic agent (anti-diabetic drug) of the new dipeptidyl peptidase-4 (DPP-4) inhibitor class of drugs. Vildagliptin inhibits the inactivation of GLP-1 and GIP by DPP-4, allowing GLP-1 and GIP to potentiate the secretion of insulin in the beta cells and suppress glucagon release by the alpha cells of the islets of Langerhans in the pancreas. Vildagliptin has been shown to reduce hyperglycemia in type 2 diabetes mellitus.

##### *tablets*

60 x 50 mg

#### AGNIS COMBI

##### VILDAGLIPTIN AND METFORMIN

Agnis Combi is indicated in the treatment of type 2 diabetes:

- in the treatment of adult patients in whom satisfactory results cannot be achieved glycemic control using oral metformin in the maximum tolerated dose or in patients who are already treated with a combination of separate vildagliptin and metformin tablets.
- in combination with a sulphonylurea (ie triple combined therapy) as an adjunct diet and exercise in adult patients inadequately controlled with metformin and sulphonylurea.
- in triple combination therapy with insulin as a supplement to diet and exercise for improving glycemic control in adult patients in whom a stable dose of insulin and metformin does not enable adequate glycemic control

##### *film-coated tablets*

60 x (50 mg / 850 mg)

60 x (50 mg / 1000 mg)

#### BELFORMIN

##### METFORMIN

##### *film-coated tablets*

100 x 500 mg

30 x 850 mg

30 x 1000 mg

120 x 850 mg

120 x 1000 mg

## BELOSEPT

### HEXETIDINE

Oral antiseptic, a broad-spectrum antibacterial and antimycotic agent for rinsing the mouth.

#### *solution*

200 ml (1 mg / ml)

## GLIKA

### GLICLAZIDE

Hypoglycemic sulphonylurea oral antidiabetic active substance. Reduces blood glucose levels by stimulating insulin secretion from  $\beta$ -cells of the islands of Langerhans in pancreas.

#### *modified-release tablets*

60 x 30 mg      30 x 60 mg  
60 x 60 mg  
90 x 60 mg

## LORDIAR

### LOPERAMIDE

Synthetic antidiarrheal for oral use.

#### *hard capsules*

20 x 2 mg

## PORTALAK efekt

### LACTULOSE

Osmotic laxative, used in occasional or long-term constipation.

#### *syrup*

250 ml (66.7 g / 100 ml)

## PORTALAK

### LACTULOSE

Osmotic laxative, used in occasional or long-term constipation.

#### *syrup*

500 ml (66.7 g / 100 ml)

## POTASSIUM CHLORIDE BELUPO (KALIJEV KLORID BELUPO)

### POTASSIUM CHLORIDE

Oral replacement of electrolyte and fluid loss due to various reasons.

#### *soluble tablets*

30 x 500 mg

## REGLINID

### REPAGLINIDE

Antidiabetic drug, an oral blood glucose-lowering drug of the meglitinide class, used in the management of non-insulin-dependent diabetes mellitus (NIDDM). Repaglinide lowers blood glucose levels by stimulating the release of insulin from the pancreas.

#### *tablets*

90 x 0.5 mg      90 x 1 mg      90 x 2 mg

## ROJAZOL

### MICONAZOLE

Topical broad-spectrum antimycotic. Combined fungistatic and fungicidal action, bacteriostatic and bactericidal action on gram-positive cocci and bacilli that often follow fungous infections

*oral gel*

40 g 2% (20 mg / g)

## RUDAKOL

### MEBEVERINE

Selective musculotropic spasmolytic without anticholinergic side effects.

*coated tablets*

50 x 135 mg

*prolonged-release hard capsules*

60 x 200 mg

## TARGET PLUS

### OMEPRAZOLE

Proton pumps inhibitor, effective in the short-term treatment of duodenal, gastric and esophageal ulcers and may offer a useful alternative to conventional therapy. In combination with antimicrobials in dual and triple drug regimens, omeprazole therapy can achieve high cure rates in the treatment of *Helicobacter pylori* associated with peptic ulcer disease. The drug is recommended for Zollinger-Ellison syndrome.

*hard capsules*

28 x 20 mg

## ZYNOL

### BENZYDAMINE

ZYNOL 1.5 mg/ml spray for the oral mucosa, solution is a locally acting analgesic and anti-inflammatory drug for painful conditions in the throat and mouth in adults and children over 6 years of age.

Multidose spray HDPE container, fitted with a metering pump and nozzle, containing 30 ml

ZYNOL 3 mg/ml spray for oral mucosa, solution is a locally acting analgesic and anti-inflammatory drug for painful conditions and swelling in the pharynx and mouth in adults.

Multidose spray HDPE container, fitted with a metering pump and nozzle, containing 15 ml

ZYNOL 1.5 mg/ml solution for gargling/rinsing the mouth is a locally acting analgesic and anti-inflammatory drug for painful conditions in the pharynx and mouth in adults and adolescents.

*Clear glass bottle with screw cap containing 120 ml, with graduated 20 ml measuring cup.*

## ZOLTEX

### PANTOPRAZOLE

Proton pumps inhibitor, effective in the short-term treatment of duodenal, gastric and esophageal ulcers and may offer a useful alternative to conventional therapy. In combination with antimicrobials in dual and triple drug regimens, omeprazole therapy can achieve high cure rate in the treatment of *Helicobacter pylori* associated with peptic ulcer disease. The drug is recommended for Zollinger-Ellison syndrome.

*gastro-resistant tablets*

28 x 20 mg

56 x 20 mg

28 x 40 mg

*powder for solution for injection*

40 mg / 10 ml



# B

## BLOOD AND BLOOD FORMING ORGANS

### ASKA pro

#### ACETYLSALICYLIC ACID

Acetylsalicylic acid inhibits the adhesion and dumping of blood platelets (thrombocytes) and in this way prevents blood clots (thrombi) to develop (inhibition of platelet aggregation).

*gastro-resistant tablets*

30 x 100 mg

### GRECCAS

#### TICAGRELOR

Greccas, co-administered with acetylsalicylic acid (ASA), is indicated for the prevention of atherothrombotic events in adult patients with acute coronary syndrome (ACS).

*film-coated tablets*

56 x 90 mg

### HEPARIN BELUPO

#### HEPARIN SODIUM

Endogenous anticoagulant for intravenous infusion or intravenous injection.

*solution for injections*

10 x 5 ml (25,000 IU / 5 ml)

### KLOPIDEX

#### CLOPIDOGREL

Clopidogrel is an oral antiplatelet agent (thienopyridine class) which inhibits blood clotting and is indicated for prevention of vascular ischaemic events in patients with symptomatic atherosclerosis, acute coronary syndrome without ST-segment elevation (NSTEMI), ST elevation MI (STEMI). It is also used, along with aspirin, for the prevention of thrombosis after placement of intracoronary stent or as an alternative antiplatelet drug for patients who are intolerant to aspirin.

*film-coated tablets*

30 x 75 mg

## CARDIOVASCULAR SYSTEM

### AMIODARON HYDROCHLORIDE BELUPO

#### AMIODARONE HYDROCHLORIDE

Antiarrhythmic drug, class III.

##### *tablets*

60 x 200 mg

### AMONEX

#### AMLODIPINE

Third generation of the dihydropyridine derivative. Calcium antagonist with predominantly vascular effect, indicated for the treatment of essential hypertension, chronic stable angina pectoris and Prinzmetal angina.

##### *tablets*

30 x 5 mg      30 x 10 mg

60 x 5 mg      60 x 10 mg

90 x 5 mg      90 x 10 mg

### AMORA

#### RAMIRPIL AND AMLODIPINE

The combination of an ACE inhibitor ramipril and a dihydropyridine calcium channel blocker amlodipine. A fixed combination of ramipril with amlodipine effectively provides both blood pressure control and positive cardioprotective and nephroprotective effect.

##### *hard capsules*

30 x 2,5 mg/5 mg      60 x 2,5 mg/5 mg

30 x 5 mg/5 mg      60 x 5 mg/5 mg

30 x 5 mg/10 mg      60 x 5 mg/10 mg

30 x 10 mg/5 mg      60 x 10 mg/5 mg

30 x 10 mg/10 mg      60 x 10 mg/10 mg

### BISOBEL

#### BISOPROLOL

Cardioselective  $\beta_1$ -adrenergic receptor blocker, indicated for treatment of essential hypertension, coronary heart disease and congestive heart failure.

##### *tablets*

30 x 2.5 mg

60 x 2.5 mg

30 x 5 mg

60 x 5 mg

30 x 10 mg

## CARVELOL

### CARVEDILOL

$\alpha$ - and  $\beta$ -adrenergic receptor blocker, indicated for the treatment of essential hypertension, coronary heart disease and symptomatic congestive heart failure.

#### *tablets*

30 x 3.125 mg  
30 x 6.25 mg  
30 x 12.5 mg  
30 x 25 mg

## DOXA

### DOXAZOSIN

Alpha-adrenergic blockers. It causes the blood vessels (veins and arteries) to relax and expand, improving blood flow. Doxazosin also relaxes muscles in the prostate and bladder neck, making it easier to urinate.

#### *tablets*

20 x 2 mg      20 x 4 mg

## FURSEMID

### FUROSEMIDE

Potent loop diuretic. Inhibits sodium and chloride reabsorption at the loop of Henle.

#### *tablets*

20 x 40 mg

#### *solution for injections*

5 x 2 ml (20 mg / 2 ml)

## FURSEMID FORTE

### FUROSEMIDE

Potent loop diuretic. Inhibits sodium and chloride reabsorption at the loop of Henle.

#### *tablets*

20 x 500 mg

## INDAMID SR

### INDAPAMIDE

Diuretic and antihypertensive. Pharmacologically related to thiazide diuretics with antihypertensive action.

#### *prolonged-release tablets*

30 x 1.5 mg

## IRUMED

### LISINOPRIL

ACE inhibitor, used in hypertension, heart failure and for short-term treatment hemodynamically stable patients after acute myocardial infarction. Slows down the development of diabetic nephropathy and retinopathy.

#### *tablets*

30 x 2.5 mg	30 x 5 mg	30 x 10 mg	30 x 20 mg
60 x 2.5 mg	60 x 5 mg	60 x 10 mg	60 x 20 mg

## IRUZID

### LISINOPRIL AND HYDROCHLOROTHIAZIDE

Combination of ACE inhibitor and diuretic, antihypertensive

#### *tablets*

30 x (10 + 12.5 mg)

30 x (20 + 12.5 mg)

30 x (20 + 25 mg)

60 x (10 + 12.5 mg)

60 x (20 + 12.5 mg)

60 x (20 + 25 mg)

## ISOCARD

### ISOSORBIDE MONONITRATE

Used in prevention of angina pectoris attacks.

#### *prolonged-release tablets*

50 x 40 mg

50 x 60 mg

## LACIDIPIN BELUPO

### LACIDIPINE

Calcium channel blocker, indicated in adults for the treatment of hypertension, either alone or in combination with other antihypertensive agents.

#### *film-coated tablets*

28 x 4 mg

28 x 6 mg

## MOKSONIDIN BELUPO

### MOKSONIDINE

Antihypertensive centrally acting antiadrenergic agent.

#### *film-coated tablets*

28 x 0.2 mg

28 x 0.4 mg

## NIBEL

### NEBIVOLOL

Antihypertensive and selective  $\beta_1$ -blocker with nitric oxide-potentiating vasodilatory effect, indicated for treatment of essential hypertension and congestive heart failure.

#### *tablets*

14 x 5 mg

28 x 5 mg

## ORMIDOL

### ATENOLOL

Antihypertensive and selective  $\beta$ -blocker

#### *film-coated tablets*

30 x 25 mg

30 x 50 mg

28 x 100 mg

## PINOX

### LERCANIDIPINE

Third generation of the dihydropyridine derivative. Calcium antagonist with predominantly vascular effect, indicated for the treatment of essential hypertension. Lercanidipine works by relaxing and opening the blood vessels allowing the blood to circulate more freely around the body. This lowers the blood pressure and allows the heart to work more efficiently.

#### *film-coated tablets*

28 x 10 mg                      56 x 10 mg

## PROTECTA

### SIMVASTATINE

Hypolipemic, an HMG-CoA reductase inhibitor.

#### *film-coated tablets*

28 x 10 mg                      28 x 20 mg                      28 x 40 mg  
56 x 20 mg

## RAMED

### RAMIPRIL

Antihypertensive. ACE inhibitor, indicated for the treatment of hypertension, heart failure and post-myocardial infarction. Slows down the development of non-diabetic and diabetic nephropathy.

#### *tablets*

28 x 1.25 mg    28 x 2.5 mg                      28 x 5 mg                      28 x 10 mg  
56 x 2.5 mg                      56 x 5 mg

## RAMZID

### RAMIPRIL AND HYDROCHLOROTHIAZIDE

Antihypertensive. Combination of ACE inhibitor and diuretic.

#### *tablets*

28 x (2.5 + 12.5 mg)                      28 x (5 + 25 mg)  
56 x (2.5 + 12.5 mg)                      56 x (5 + 25 mg)

## ROSIX

### ROSUVASTATINE

Hypolipemic, an inhibitor of HMG-CoA reductase, reduces cholesterol by increasing the number of low-density lipoprotein (LDL) receptors on the cell-surface to enhance uptake and catabolism of LDL, inhibits hepatic synthesis of hepatic very-low-density lipoprotein (VLDL), which reduces the total number of VLDL and LDL particles. The treatment reduces triglycerides (TG) and produces increases in high-density lipoprotein cholesterol (HDL).

#### *film-coated tablets*

28 x 10 mg  
28 x 20 mg  
28 x 40 mg

## ROSIX AM

### ROSUVASTATINE AND AMLODIPINE

Rosuvastatine/amlodipine is indicated as substitution therapy for those patients who are adequately controlled with rosuvastatin and amlodipine given concurrently, at the same dose level as in the combination for the treatment of hypertension in adult patients who are estimated to have a high risk for a first cardiovascular event (for prevention of major cardiovascular events) as an adjunct to correction of other risk factors or with one of the following coincident conditions:

- primary hypercholesterolaemia (type IIa including heterozygous familial hypercholesterolaemia) or mixed dyslipidaemia (type IIb) as an adjunct to diet when response to diet and other non-pharmacological treatments (eg exercise, weight reduction) is inadequate
- homozygous familial hypercholesterolaemia as an adjunct to diet and other lipid lowering treatments (eg LDL apheresis) or if such treatments are not appropriate.

#### *hard capsules*

30 x (10 mg / 5 mg)                      30 x (10 mg / 10 mg)

30 x (20 mg / 5 mg)                      30 x 20 mg / 10 mg)

## ROSIX COMBI

### ROSUVASTATIN AND EZETIMIBE

Rosuvastatine/ezetimibe is indicated as adjunct to diet for treatment of primary hypercholesterolemia as substitution therapy in adult patients adequately controlled with the individual substances given concurrently at the same dose level as in the fixed dose combination, but as separate products.

#### *tablets*

30 x (5 mg / 10 mg)

30 x (10 mg / 10 mg)

30 x (20 mg / 10 mg)

## ROSIX DUO

### ROSUVASTATIN AND ACETYLSALYSLIC ACID

ROSIX Duo is indicated for the secondary prevention of cardiovascular events as a replacement therapy in adult patients who are adequately controlled with monocomponents given simultaneously in equivalent therapeutic doses.

#### *hard capsules*

30 x (5 mg / 100 mg)

30 x (10 mg / 100 mg)

30 x (20 mg / 100 mg)

## TOMID

### TORASEMIDE

Diuretic and antihypertensive. Torasemide is a pyridine-sulfonyl urea type loop diuretic mainly used in the management of edema associated with congestive heart failure. It is also used at low doses for the management of hypertension.

#### *tablets*

30 x 5 mg

30 x 10 mg



## TORVAS

### ATORVASTATINE

Hypolipemic. Indicated as an adjunct to diet for reduction of elevated total cholesterol, LDL-cholesterol, apolipoprotein B, and triglycerides in patients with primary hypercholesterolaemia, heterozygous familial hypercholesterolaemia or combined (mixed) hyperlipidaemia when response to diet and other nonpharmacological measures is inadequate.

#### *film-coated tablets*

30 x 10 mg	30 x 20 mg	30 x 40 mg
60 x 10 mg	60 x 20 mg	60 x 40 mg

## VAINAS

### AMLODIPINE AND VALSARTANE

Treatment of essential hypertension, in adults whose blood pressure is not adequately controlled on amlodipine or valsartan monotherapy.

#### *film-coated tablets*

30 x (5 mg / 80 mg)  
30 x (5 mg / 160 mg)  
30 x (10 mg / 160 mg)

## VAINAS HCT

### AMLODIPINE AND VALSARTANE AND HYDROCHLOROTHIAZIDE

Treatment of essential hypertension as substitution therapy in adult patients whose blood pressure is adequately controlled on the combination amlodipine, valsartan and hydrochlorothiazide (HCT), taken either as three single-component formulations or as a dual-component and a single-component formulation.

#### *film-coated tablets*

30 x (5 mg / 160 mg / 12,5 mg)	30 x (10 mg / 160 mg / 12,5 mg)
30 x (5 mg / 160 mg / 25 mg)	30 x (10 mg / 160 mg / 25 mg)

## VAL

### VALSARTAN

Antihypertensive. Angiotensin II antagonist acting on the AT1 receptor, indicated for the treatment of hypertension, heart failure and post-myocardial infarction.

#### *film-coated tablets*

28 x 80 mg	28 x 160 mg
56 x 80 mg	56 x 160 mg

## VAL PLUS

### VALSARTAN AND HYDROCHLOROTHIAZIDE

Antihypertensive. Combination of Angiotensin II antagonist and diuretic.

#### *film-coated tablets*

28 x (80 + 12.5 mg)	28 x (160 + 12.5 mg)
56 x (80 + 12.5 mg)	56 x (160 + 12.5 mg)

# D

## DERMATOLOGICALS

### AFLODERM

#### ALCLOMETHASONE DIPROPIONATE

Non-fluorinated corticosteroid dermatological with antiinflammatory, antiallergic and vasoconstrictive action, suitable for use in all age groups.

##### *cream*

20 g (0.5 mg / g)  
40 g (0.5 mg / g)

##### *ointment*

20 g (0.5 mg / g)

### ALOMAX

#### MINOXIDIL

Treatment of androgenic alopecia in men and women between 18 and 65 years. Alomax solution works as a vasodilator and therefore improves blood circulation in the scalp.

##### *solution*

60 ml 5% (for men only)  
60 ml 2% (for men and women)

### AMOFIN

#### AMOROLFINE

Broad spectrum antimycotic for topical use. Used for treatment of onychomycosis without matrix involvement, caused by dermatophytes, yeasts or moulds.

##### *medicated nail lacquer*

2.5 ml (50 mg / ml)  
the pack contains 30 cleansing swabs,  
10 spatulas and 30 nail files

5 ml (50 mg / ml)  
the pack contains 30 cleansing swabs,  
10 spatulas and 60 nail files

### BELODERM

#### BETAMETHASONE DIPROPIONATE

Fluorinated corticosteroid dermatological with antiinflammatory, antipruriginous, antiallergic and vasoconstrictive action, suitable for treatment of chronic and persistent dermatoses.

##### *cream*

15 g (0.5 mg / g)  
30 g (0.5 mg / g)  
100 g (0.5 mg / g)

##### *ointment*

15 g (0.5 mg / g)  
30 g (0.5 mg / g)

##### *skin solution*

50 ml (0.5 mg / g)

### BELODIN A Derm

#### DIMETINDENMALEAT

Dimetindene is an antiallergic drug. Dimetindene is an antihistamine/anticholinergic that is a selective H1 antagonist.

##### *gel*

30 g (1 mg/1 g)

## BELOGENT

### BETAMETHASONE DIPROPIONATE AND GENTAMICIN SULPHATE

Fluorinated corticosteroid dermatological combined with a wide-spectrum antibiotic, suitable for treatment of primary and secondary infected dermatoses.

#### *cream*

15 g (0.5 + 1 mg / g)

#### *ointment*

15 g (0.5 + 1 mg / g)

## BELOSALIC

### BETAMETHASONE DIPROPIONATE AND SALICYLIC ACID

Fluorinated corticosteroid dermatological with keratolytic, suitable for treatment of dry, peeling and hyperkeratotic dermatoses.

#### *ointment*

30 g (0.5 + 30 mg / g)

#### *skin solution*

50 ml (0.5 + 20 mg / g)

100 ml (0.5 + 20 mg / g)

## HERPLEX

### ACYCLOVIR

Highly selective antiviral for treatment of infections caused by herpes simplex and varicella-zoster viruses.

#### *cream*

2 g (50 mg / g)

## HERPLEXIM

### ACYCLOVIR

Highly selective antiviral for treatment of infections caused by herpes simplex and varicella-zoster viruses.

#### *cream*

5 g (50 mg / g)

10 g (50 mg / g)

## MIROBACT

### MUPIROCIN

Mupirocin is used as a topical treatment for primary and secondary bacterial skin infections.

#### *ointment*

15 g 2% (20 mg / g)

## ROJAZOL

### MICONAZOLE NITRATE

Topical broad-spectrum antimycotic. Combined fungistatic and fungicidal action, and bacteristatic and bactericidal action on gram-positive cocci and bacilli that often follow fungous infections.

#### *cream*

30 g 2% (20 mg / g)

## ROJAZOLDERM

### MICONAZOLE NITRATE

Topical treatment for infections on the skin, nails, hands, feet, outer ears, trunk, or scalp. The active ingredient miconazole destroys fungi and bacteria that cause infection.

#### *cream*

20 g (20 mg/g)

## SONA

### ADAPALENE

Topical retinoid suitable for treatment of mild-moderate acne.

*gel*

30 g (1 mg / g)

*cream*

30 g (1 mg / g)

## SONA DUO

### ADAPALENE AND BENZOYL PEROXIDE

Topical treatment of Acne vulgaris, when comedones, numerous papules and pustules are present. SONA DUO is indicated in adults, adolescents and children aged 9 years and older.

*gel*

30 g (1 mg / g + 25 mg / g)

## TERBINAX

### TERBINAFINE

Antimycotic. Terbinafine is used to treat certain types of fungal infection. It works by stopping the growth of fungus.

*tablets*

14 x 250 mg    42 x 250 mg

G

## GENITO URINARY SYSTEM AND SEX HORMONES

## BELOFLUX

### SOLIFENACIN

Symptomatic treatment of urge incontinence and / or increased urinary frequency and urgency as may occur in patients with overactive bladder syndrome.

*film-coated tablets*

30 x 5 mg

30 x 10 mg

## KLLOTAS

### DUTASTERIDE AND TAMSULOSIN

Treatment of moderate to severe symptoms of benign prostatic hyperplasia (BPH). Reduction in the risk of acute urinary retention (AUR) and surgery in patients with moderate to severe symptoms of BPH.

*hard capsules*

30 x (0,5 mg / 0,4 mg)

## MEDAZOL

### METRONIDAZOLE

Antiinfective, antiprotozoal used for treatment of anaerobic infections, protozoal diseases and for prophylaxis in surgical and gynecological procedures.

*vaginal tablets*

10 x 500 mg

## ROJAZOL

### MICONAZOLE NITRATE

Topical broad-spectrum antimycotic. Combined fungistatic and fungicidal action, bacteriostatic and bactericidal action on gram-positive cocci and bacilli that often follow fungous infections.

*vaginal suppositories*

7 x 200 mg

## TAMOSIN

### TAMSULOSIN

An antiadrenergic agent, is used to treat enlargement of the prostate gland in men, a condition called benign prostatic hyperplasia or BPH.

*hard prolonged-release capsules*

30 x 0.4 mg

100 x 0.4 mg

## VINER

### SILDENAFIL

In the presence of sexual stimulation, sildenafil enhances the effect of nitric oxide (NO) by inhibiting phosphodiesterase type 5 (PDE5), which is responsible for degradation of cyclic guanosine monophosphate (cGMP) in the corpus cavernosum.

*film-coated tablets*

4 x 50 mg

4 x 100 mg

## GENERAL ANTIINFECTIVES FOR SYSTEMIC USE

## AMOKSICILIN BELUPO

AMOXICILLIN TRIHYDRATE

Semi-synthetic penicillin wide-spectrum antibiotic.

*tablets for oral suspension*

16 x 500 mg

16 x 1000 mg

## AZITROMICIN BELUPO

AZITHROMYCIN

Semi-synthetic macrolide antibiotic.

*film-coated tablets*

6 x 250 mg

3 x 500 mg

## BELOXIM

CEFUROXIME

Antibacterial for systemic use in adults and children, second-generation cephalosporin.

*film-coated tablets*

10 x 250 mg

10 x 500 mg

## CEFALEKSIN BELUPO

CEFALEXIN

Semi-synthetic cephalosporin bactericidal antibiotic.

*hard capsules*

16 x 500 mg

*powder for oral suspension*

100 ml (250 mg / 5 ml)

## CIFLOX

CIPROFLOXACIN

Synthetic antibiotic belong to a group of fluoroquinolones. Broad-spectrum antibiotic that is active against both Gram-positive and Gram-negative bacteria.

*film-coated tablets*

10 x 250 mg

10 x 500 mg



## DOKSICIKLIN BELUPO

### DOXYCYCLINE HYCLATE

Tetracycline wide-spectrum antibiotic.

*hard capsules*

6 x 100 mg      25 x 100 mg

## ERITROMICIN BELUPO

### ERYTHROMYCIN ESTOLATE

Macrolide wide-spectrum antibiotic.

*hard capsules*

16 x 250 mg

## GENTAMICIN BELUPO

### GENTAMICIN SULPHATE

Aminoglycoside wide-spectrum antibiotic, for intramuscular and intravenous use.

*solution for injections*

10 x 2 ml (80 mg / 2 ml)      10 x 2 ml (120 mg / 2 ml)

## HERPLEXIM

### ACYCLOVIR

Highly selective antiviral for treatment of infections caused by herpes simplex and varicella-zoster viruses.

*film-coated tablets*

35 x 400 mg

## ISOTRETINOIN BELUPO

### ISOTRETINOIN

Broad-spectrum oral antimycotic.

*soft gel capsules*

30 x 10 mg      30 x 20 mg  
60 x 10 mg      60 x 20 mg

## ITRAC

### ITRACONAZOLE

Retinoid for treatment of acne. Indicated for severe forms of acne (such as nodular or conglobate acne or acne at risk of permanent scarring) resistant to adequate courses of standard therapy with systemic anti-bacterials and topical therapy.

*hard capsules*

4 x 100 mg      28 x 100 mg

## KLAVOBEL BID

### AMOXICILIN TRIHYDRATE + POTASSIUM CLAVULANATE

Semi-synthetic penicillin antibiotic used to treat a wide range of infections caused by bacteria.

*film-coated tablets*

14 x (875 + 125) mg

## MAKCIN

### CLARITHROMYCIN

Broad-spectrum macrolide antibiotic.

*film-coated tablets*

14 x 500 mg

## MAKCIN SR

### CLARITHROMYCIN

Broad-spectrum macrolide antibiotic.

*prolonged-release tablets*

7 x 500 mg

## NINUR

### NITROFURANTOIN

An urinary anti-infective, used to treat specific urinary tract infections.

*hard capsules*

30 x 50 mg

## NOFLOX

### NORFLOXACIN

Broad-spectrum, quinolone bactericidal agent indicated for the treatment of upper and lower, complicated and uncomplicated, acute and chronic urinary tract infections.

*film-coated tablets*

20 x 400 mg

## SILAPEN K

### PHENOXYMETHYLPENICILLIN POTASSIUM

Semi-synthetic penicillin antibiotic, particularly suitable for treatment of streptococcal infections.

Each 5 ml of oral solution contains 250 mg of Phenoxymethylpenicillin as Phenoxymethylpenicillin Potassium Ph. Eur.

*powder for oral solution*

100 ml (250 mg / 5 ml)

## SULOTRIM

### SULFAMETHOXAZOLE AND TRIMETHOPRIM

Combined antibacterial agent having two active components with a synergetic effect.

*tablets*

20 x (400 + 80 mg)

## SULOTRIM FORTE

### SULFAMETHOXAZOLE AND TRIMETHOPRIM

Combined antibacterial agent having two active components with a synergetic effect.

*tablets*

20 x (800 + 160 mg)

## ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS

### BIXALAN

#### BICALUTAMID

Antineoplastic agent, antiandrogen, non-steroidal androgen receptor inhibitor. It competitively inhibits the action of androgens by binding to cytosol androgen receptors in the target tissue.

*film-coated tablets*

28 x 50 mg

28 x 150 mg

### LETRILAN

#### LETROZOLE

Nonsteroidal competitive inhibitor of the aromatase enzyme system; it inhibits the conversion of androgens to estrogens. Letrozole inhibits the aromatase enzyme by competitively binding to the heme of the cytochrome P450 subunit of the enzyme, resulting in a reduction of estrogen biosynthesis in all tissues.

*film-coated tablets*

30 x 2.5 mg

### STRAZOLAN

#### ANASTROZOLE

Potent and highly selective nonsteroidal aromatase inhibitor. Anastrozole selectively inhibits the conversion of androgens to estrogens by inhibiting the aromatase enzyme complex through competitive binding to the haeme of the cytochrome P450 (CYP450) unit of the enzyme. Suppression of estrogen biosynthesis in all tissues reduces serum concentrations of circulating estrogens, including estrone, estradiol, and estrone sulphate with a subsequent beneficial effect in breast cancer.

*film-coated tablets*

28 x 1 mg

## MUSCULO-SKELETAL SYSTEM

### ABUXAR 80 mg

#### FEBUXOSTAT

Febuxostat is indicated for the treatment of chronic hyperuricaemia in conditions where urate deposition has already occurred (including a history, or presence of, tophus and/or gouty arthritis). Febuxostat Tablets are indicated in adults.

*film-coated tablets*

28 x 80 mg

### ABUXAR 120 mg

#### FEBUXOSTAT

Febuxostat is indicated for the treatment of chronic hyperuricaemia in conditions where urate deposition has already occurred (including a history, or presence of, tophus and/or gouty arthritis).

Febuxostat is indicated for the prevention and treatment of hyperuricaemia in adult patients undergoing chemotherapy for haematologic malignancies at intermediate to high risk of Tumor Lysis Syndrome (TLS).

Febuxostat is indicated in adults.

*film-coated tablets*

28 x 120 mg

### ALOPURINOL BELUPO

#### ALLOPURINOL

An antigout agent. The primary use of allopurinol is to treat hyperuricemia and its complications. Allopurinol does not alleviate acute attacks of gout, but is useful in chronic gout to prevent future attacks.

*tablets*

100 x 100 mg

30 x 200 mg

100 x 200 mg

### BONNA

#### RISEDRONATE SODIUM

Risedronate is a potent bisphosphonate in the treatment of established postmenopausal osteoporosis in the women, osteoporosis in the man and osteoporosis caused by medicines (glucocorticoides).

*film-coated tablets*

4 x 35 mg

2 x 75 mg

6 x 75 mg

## IBUPROFEN BELUPO

### IBUPROFEN

Non-steroidal anti-inflammatory agent (NSAID) with analgesic and antipyretic effect. Rapidly and efficiently removes severe, moderate and sudden pain in acute rheumatic conditions, in soft tissue injuries, acute disorders of the lumbar spinal column and after surgery.

#### *film-coated tablets*

30 x 400 mg

30 x 600 mg

30 x 800 mg

#### *oral suspension*

100 ml (100 mg / 5 ml)

## INDOMETACIN BELUPO

### INDOMETACIN

Non-steroidal anti-inflammatory agent (NSAID). Helps to decrease swelling and pain.

#### *hard capsules*

30 x 25 mg

## KNAVON

### KETOPROFEN

Non-steroidal anti-inflammatory agent (NSAID). Used in inflammatory and degenerative rheumatic diseases to reduce joint swelling, to reduce pain both at rest and in motion, to shorten the duration of morning stiffness, enable greater flexibility and functional capacity. Also suitable for removing abdominal pain.

#### *hard capsules*

25 x 50 mg

#### *suppositories*

12 x 100 mg

## KNAVON FORTE

### KETOPROFEN

Non-steroidal anti-inflammatory agent (NSAID). Used in inflammatory and degenerative rheumatic diseases to reduce joint swelling, to reduce pain both at rest and in motion, to shorten the duration of morning stiffness, enable greater flexibility and functional capacity. Also suitable for removing abdominal pain.

#### *film-coated tablets*

20 x 100 mg

## KNAVON RETARD

### KETOPROFEN

Non-steroidal anti-inflammatory agent (NSAID). Used in inflammatory and degenerative rheumatic diseases to reduce joint swelling, to reduce pain both at rest and in motion, to shorten the duration of morning stiffness, enable greater flexibility and functional capacity. Also suitable for removing abdominal pain.

#### *prolonged-release tablets*

20 x 150 mg

## KNAVON AKUT

### KETOPROFEN

KNAVON AKUT 50 mg hard capsules are indicated for short-term symptomatic relief of mild to moderate acute pain such as: headaches, toothache, menstrual pains, pain after minor sprains and dislocations.

For the treatment of adults and adolescents over 16 years of age.

#### *hard capsules*

10 x 50 mg

## LUBOR

### PIROXICAM

Non-steroidal anti-inflammatory agent (NSAID), which effectively removes edema, erythema, and tissue proliferation in the affected joint. Used as analgesic in supporting therapy.

#### *hard capsules*

20 x 10 mg    20 x 20 mg

#### *suppositories*

10 x 20 mg

## LUBORETA

### PIROXICAM

A topical NSAID that relieves pain due to sprains, contusions, strained muscles and for treatment of particular rheumatism.

#### *cream*

50 g (10 mg / g)

## NEOFEN 200 mg tablets

### IBUPROFEN

Non-steroidal anti-inflammatory agent (NSAID) with analgesic and antipyretic effect. Removes mild and moderate pain of various origin (headache, toothache, premenstrual and menstrual pain, pain caused by injury, rheumatic diseases, cold and influenza). Particularly suitable for treatment of persistent high temperature (over 39°C).

#### *film-coated tablets*

20 x 200 mg

30 x 200 mg

## NEOFEN RAPID

### IBUPROFEN LYSINE

NEOFEN RAPID is indicated in the treatment of the following conditions: mild to moderately painful conditions (eg back pain, toothache, muscle pain, joint pain, menstrual pain, neuropathic pain); acute phase of migraine with or without aura and for the treatment of tension headaches; pain due to common cold and flu.

NEOFEN RAPID is intended for use in adults and adolescents over 12 years of age (body weight  $\geq$  40 kg).

#### *film-coated tablets*

10 x 342 mg

10 x 684 mg

## NEOFEN forte

### IBUPROFEN

Non-steroidal anti-inflammatory agent (NSAID) with analgesic and antipyretic effect. Removes mild and moderate pain of various origin (headache, toothache, premenstrual and menstrual pain, pain caused by injury, rheumatic diseases, cold and influenza). Particularly suitable for treatment of persistent high temperature (over 39°C).

#### *film-coated tablets*

10 x 400 mg

20 x 400 mg



## NEOFEN NEO forte

### IBUPROFEN

NEOFEN NEO FORTE is intended for use in adult patients and adolescents over 12 years of age (body weight greater than 40 kg) for short-term symptomatic treatment: mild to moderate pain such as headache (including migraine), neuralgia, toothache, back pain, muscle and joint pain, and menstrual pain; reduction of elevated body temperature and relief of pain and discomfort due to common cold and flu.

Shaped like a capsule, caplet serves as an easy-to-swallow alternative to regular tablets. Does not contain lactose or starch.

#### *caplets*

20 x 400 mg

## NEOFEN plus gel

### IBUPROFEN

Non-steroidal anti-inflammatory agent (NSAID) for topical treatment. Relieves pain due to sprains, contusions, strained muscles and for treatment of particular rheumatism and sports injuries.

#### *gel*

50 g (50 mg / g)

100 g (50 mg / g)

## NEOFEN oral suspension

### IBUPROFEN

Non-steroidal anti-inflammatory agent (NSAID). It is used for treatment of mild to moderate pain, post-immunization pyrexia, rheumatic or muscular pain, headache, reduction of fever, sore throat, teething pain, toothache, minor aches and pains, symptoms of cold and influenza.

#### *oral suspension*

100 ml (100 mg / 5 ml)

## NEOFEN 60 g suppositories

### IBUPROFEN

Non-steroidal anti-inflammatory agent (NSAID). It is used for treatment of mild to moderate pain, post-immunization pyrexia, rheumatic or muscular pain, headache, reduction of fever, sore throat, teething pain, toothache, minor aches and pains, symptoms of cold and influenza in children age 3 months to 2 years.

#### *suppositories*

10 x 60 mg

## NEOFEN 125 g suppositories

### IBUPROFEN

Non-steroidal anti-inflammatory agent (NSAID). It is used for treatment of mild to moderate pain, post-immunization pyrexia, rheumatic or muscular pain, headache, reduction of fever, sore throat, teething pain, toothache, minor aches and pains, symptoms of cold and influenza in children age 2-6 years.

#### *suppositories*

10 x 125 mg



## NEOFEN Combo

### IBUPROFEN AND ACETAMINOPHEN (PARACETAMOL)

For short-term relief of mild to moderate pain associated with migraine, headache, back pain, menstrual pain, toothache, rheumatic and muscle pain, non-inflammatory arthritis pain, as well as cold and flu symptoms, sore throat and fever. This medicine is particularly suitable for pain that requires stronger analgesia than with ibuprofen or paracetamol alone.

NEOFEN COMBO is indicated for adults over 18 years of age.

#### *film-coated tablets*

10 x (200 mg / 500 mg)

20 x (200 mg / 500 mg)

## N

## NERVOUS SYSTEM



## ARVIND

### LAMOTRIGINE

Anticonvulsant. Used alone or in combination with other medications in the treatment of seizures in adults and children with epilepsy and Lennox-Gastaut syndrome. Lamotrigine is also used in the treatment of bipolar disorder.

#### *tablets*

30 x 25 mg

30 x 50 mg

30 x 100 mg



## AZOLAR

### ARIPRAZOLE

Antipsychotic indicated for the treatment schizophrenia and manic episodes in bipolar I. disorder.

#### *tablets*

30 x 10 mg

30 x 15 mg

30 x 30 mg



## CALIXTA

### MIRTAZAPINE

Tetracyclic antidepressant. Noradrenergic and specific serotonergic antidepressant (NaSSA).

#### *film-coated tablets*

30 x 15 mg

30 x 30 mg

30 x 45 mg

## CERSON

### NITRAZEPAM

Benzodiazepine hypnotic, indicated for short-term treatment of insomnia.

#### *tablets*

10 x 5 mg

30 x 5 mg

## CITRAM

### ESCITALOPRAM

An antidepressant of the selective serotonin reuptake inhibitor (SSRI) class, noted for its high selectivity of serotonin reuptake inhibition. It is the S-stereoisomer (enantiomer) of the earlier drug citalopram. It is indicated for the treatment in adults with major depressive disorder, generalized anxiety disorder and social anxiety disorder.

#### *film-coated tablets*

28 x 10 mg

## DOLYXAN

### DULOXETINE

Duloxetine is a combined serotonin (5-HT) and noradrenaline (NA) reuptake inhibitor.

It is used in the treatment of major depressive disorder, the treatment of pain in diabetic peripheral neuropathy and the treatment of generalized anxiety disorder. Dolyxan is indicated in adults.

#### *gastro-resistant capsules, hard*

28 x 30 mg

28 x 60 mg

## EMINENS SR

### ROPINIROLE

Non-ergoline D2/D3 dopamine agonist that alleviates dopamine deficiency by stimulating striatal dopamine receptors.

#### *prolonged-release tablets*

28 x 2 mg

28 x 4 mg

28 x 8 mg

## KATENA

### GABAPENTIN

A gamma-aminobutyric acid analog. Gabapentin is indicated as therapy in the treatment of partial seizures in adults with epilepsy with and without secondary generalization.

#### *capsules*

20 x 100 mg

50 x 300 mg

50 x 400 mg

## LEVETIRACETAM BELUPO

### LEVETIRACETAM

Antiepileptic, indicated as monotherapy for adults and adolescents from 16 years of age or as add-on therapy for adults ( $\geq 18$  years) and adolescents (12 to 17 years).

#### *film-coated tablets*

60 x 250 mg

60 x 500 mg

60 x 1000 mg

## LIDOKAIN BELUPO

### LIDOCAINE

A local anesthetic and antiarrhythmic

#### *solution for injections*

5 x 5 ml (2%)

100 x 2 ml (2%)

#### *spray*

50 ml 10% (100 mg / ml)

50 ml 10% (100 mg / ml)

## LIDOKAIN-ADRENALIN BELUPO

### LIDOCAINE, ADRENALIN

A local anesthetic and antiarrhythmic

#### *solution for injections*

100 x 2 ml (40 + 0.025 mg / 2 ml)

## LORSILAN

### LORAZEPAM

Benzodiazepine anxiolytic. Due to simple inactivation (no intermediate products) it is suitable for treatment of patients with liver disease and older patients.

#### *tablets*

20 x 2.5 mg

30 x 1 mg

## LUMIDOL

### TRAMADOL

Opioidic analgesic for treatment of moderate to severe acute and chronic pain.

#### *hard capsules*

20 x 50 mg

#### *oral drops*

10 ml (100 mg / ml)

#### *solution for injections*

5 x 1 ml (50 mg / ml)

5 x 2 ml (100 mg / 2 ml)

## LUMIDOL RETARD

### TRAMADOL

Opioidic analgesic for treatment of moderate to severe acute and chronic pain.

#### *prolonged-release tablets*

30 x 100 mg

30 x 200 mg

## LUPOCET 500 mg tablets

### ACETAMINOPHEN (PARACETAMOL)

Paracetamol is an antipyretic and analgesic. It is recommended to relieve pain and to reduce fever. It is particularly suitable for the treatment of patients sensitive to salicylates.

#### *tablets*

20 x 500 mg

## ● LUPOCET 500 mg film-coated tablets

### ACETAMINOPHEN (PARACETAMOL)

Paracetamol is an antipyretic and analgesic. It is recommended to relieve pain and to reduce fever. It is particularly suitable for the treatment of patients sensitive to salicylates.

#### *film-coated tablets*

18 x 500 mg

## ● LUPOCET forte 1000 mg film-coated tablets

### ACETAMINOPHEN (PARACETAMOL)

Paracetamol is an antipyretic and analgesic. It is recommended to relieve pain and to reduce fever. It is particularly suitable for the treatment of patients sensitive to salicylates.

#### *film-coated tablets*

10 x 1000 mg

## ● LUPOCET baby 120 mg suppositories

### ACETAMINOPHEN (PARACETAMOL)

LUPOCET suppositories are indicated for children for reducing fever and relieving pain caused by dentition, toothache, headache, pain after surgery or injuries, muscle and joint pain.

The use of paracetamol in suppository form is particularly suitable for patients who cannot take the oral form (e.g. after surgery or due to nausea or vomiting).

#### *suppositories*

10 x 120 mg

## ● LUPOCET 250 mg suppositories

### ACETAMINOPHEN (PARACETAMOL)

LUPOCET suppositories are indicated for children for reducing fever and relieving pain caused by dentition, toothache, headache, pain after surgery or injuries, muscle and joint pain.

The use of paracetamol in suppository form is particularly suitable for patients who cannot take the oral form (e.g. after surgery or due to nausea or vomiting).

#### *suppositories*

10 x 250 mg

## ● LUPOCET 1000 mg suppositories

### ACETAMINOPHEN (PARACETAMOL)

LUPOCET 1000 mg suppositories are indicated for the symptomatic treatment of mild to moderately severe pain and/or elevated body temperature.

The use of paracetamol in the form of suppositories is particularly suitable for patients who cannot take oral forms.

The drug is intended for use in adults and adolescents over 15 years of age whose body weight is greater than 50 kg.

#### *suppositories*

8 x 1000 mg

## ● LUPOCET flu effervescent tablets

ACETAMINOPHEN (PARACETAMOL), CHLORPHENIRAMINE MALEATE, VITAMINE C

LUPOCET FLU relieves the symptoms of cold and flu such as running and itchy nose, running eyes, hoarseness, cough, headache, fever and fatigue.

1 effervescent tablet contains 300 mg acetaminophen (paracetamol), 1.41 mg chlorpheniramine and 250 mg ascorbic acid.

*effervescent tablets*

10 x (300 mg + 1.41 mg + 250 mg)

## ● LUPOCET flu forte effervescent tablets

ACETAMINOPHEN (PARACETAMOL), CHLORPHENIRAMINE MALEATE, VITAMINE C

LUPOCET FLU forte relieves the symptoms of cold and flu such as running and itchy nose, running eyes, hoarseness, cough, headache, fever and fatigue.

1 effervescent tablet contains 500 mg acetaminophen (paracetamol), 1.41 mg chlorpheniramine and 250 mg ascorbic acid.

*effervescent tablets*

10 x (500 mg + 1.41 mg + 250 mg)

## ● LUPOCET junior syrup

ACETAMINOPHEN (PARACETAMOL)

Paracetamol is indicated for children for lowering fever associated with common cold, flu, infectious childhood diseases (e.g. chicken pox, whooping cough, measles, mumps) or reactions to vaccinations, and for relieving toothache, headache, sore throat, dentition pain, pain after surgery or injuries, muscle and joint pain.

5 ml syrup contains 120 mg acetaminophen (paracetamol).

*syrup*

100 ml

## ● LUPOCET DIREKT 500 mg granules

ACETAMINOPHEN (PARACETAMOL)

In adults, the use of paracetamol is recommended for the relief of mild to moderate pain (e.g. headache (including migraine and tension-type headache), toothache, neuralgia, back pain, rheumatic and muscle pain, dysmenorrhea and sore throat. It is recommended for the symptomatic relief of pain in mild osteoarthritis which has been previously diagnosed by a doctor. It is used to reduce elevated body temperature due to the common cold or flu.

In children, paracetamol is indicated for lowering elevated body temperature associated with colds, flu, childhood infectious diseases (e.g. chicken pox, whooping cough, measles, mumps) or a reaction to vaccination, as well as for the relief of mild to moderate pain associated with teething, toothache, headache and sore throat.

With prior consultation with a doctor, it is used after operations and injuries, as well as pain in muscles and joints.

LUPOCET DIREKT 500 mg granules are intended for short-term use in adults and adolescents over 12 years of age with a body weight of more than 40 kg. 5 ml syrup contains 120 mg acetaminophen (paracetamol).

*sachets*

10 x 500 mg



## LUPOCET teen capsules

### ACETAMINOPHEN (PARACETAMOL)

Paracetamol is an antipyretic and analgesic. Its use is recommended to relieve pain and to reduce fever in the common cold or flu, and for other pains such as headache (including migraine and tension headache), toothache, neuralgia, back pain, rheumatic and muscle pain, dysmenorrhoea and sore throat. It is also recommended for symptomatic relief of pain in mild arthritis. It is particularly suitable for the treatment of patients sensitive to salicylates.

#### *capsules*

20 x 300 mg

## MIDZA

### LACOSAMIDE

Lacosamide is an anticonvulsant compound indicated as monotherapy and adjunctive therapy in the treatment of partial-onset seizures with or without secondary generalisation in adults, adolescents and children from 4 years of age with epilepsy.

#### *film coated tablets*

14 x 50 mg

56 x 50 mg

56 x 100 mg

56 x 150 mg

56 x 200 mg

## MISAR

### ALPRAZOLAM

Benzodiazepine anxiolytic used to relieve anxiety, nervousness, and tension associated with anxiety disorders. Also used to treat panic disorders.

#### *tablets*

30 x 0.25 mg      30 x 0.5 mg      30 x 1 mg

## MISAR SR

### ALPRAZOLAM

Benzodiazepine anxiolytic (see previous).

#### *prolonged-release tablets*

30 x 0.5 mg

30 x 1 mg

## NORMABEL

### DIAZEPAM

Benzodiazepine anxiolytic, sedative and central miorelaxant. Used to relieve anxiety, nervousness, and tension associated with anxiety disorders. It is also used to treat certain types of seizure disorders and muscle spasms.

#### *film-coated tablets*

30 x 2 mg

30 x 5 mg

#### *tablets*

30 x 10 mg

#### *solution for injections*

10 x 2 ml (10 mg / 2 ml)

## OKSAZEPAM BELUPO

### OXAZEPAM

Benzodiazepine anxiolytic

#### *tablets*

30 x 10 mg    30 x 15 mg    30 x 30 mg

## PREGABALIN BELUPO

### PREGABALIN

Pregabalin is indicated for the treatment of peripheral and central neuropathic pain in adults, excluding the treatment of trigeminal neuralgia pain, acute herpetic pain, post-herpetic pain, or causalgia pain. It is indicated as adjunctive therapy in adults with partial seizures with or without secondary generalisation. Also, it is indicated for the treatment of Generalised Anxiety Disorder (GAD) in adults.

#### *coated tablets*

56 x 75 mg

56 x 150 mg

56 x 300 mg

## PROMAZIN BELUPO

### PROMAZINE

Phenothiazine class of antipsychotics. Used as an adjunct to short-term management of moderate to severe psychomotor agitation. It is also used to treat agitation and restlessness in the elderly.

#### *coated tablets*

50 x 25 mg

50 x 100 mg

## PROSPERA

### RISPERIDONE

Antipsychotic for the treatment of schizophrenia and mania associated with bipolar disorder.

#### *film-coated tablets*

20 x 1 mg

60 x 2 mg

## Q-PIN SR

### QUETIAPINE

- Treatment of schizophrenia
- Treatment of bipolar disorder
  - for the treatment of moderate to severe manic episodes in bipolar disorder
  - for the treatment of major depressive episodes in bipolar disorder
  - to prevent the recurrence of manic or depressive episodes in patients with bipolar disorder who have previously responded to treatment with quetiapine.
- Additional therapy of major depressive episodes in patients with major depressive disorder (MDD) who had a suboptimal response to antidepressant monotherapy. Before starting treatment, clinicians must consider the safety profile of Q-PIN SR

#### *prolonged-release tablets*

60 x 50 mg

60 x 150 mg

60 x 200 mg

60 x 300 mg

60 x 400 mg

## Q-PIN

### QUETIAPINE

Antipsycgotic indicated for treatment of schizophrenia and for treatment of bipolar disorder in the case of moderate to severe manic episodes, major depressive episodes in bipolar disorder and for the prevention of recurrence of manic or depressed episodes in patients with bipolar disorder who previously responded to quetiapine treatment.

#### *film-coated tablets*

60 x 25 mg

60 x 100 mg

60 x 200 mg

## SULPIRID BELUPO

### SULPIRIDE

Sulpiride is used in the treatment of:

- acute and chronic schizophrenia
- anxiety and depressive disorders

#### *hard capsules*

30 x 50 mg

12 x 200 mg

## TIRAMAT

### TOPIRAMATE

A carbonic anhydrase inhibitor anticonvulsant. It is used to treat seizures in adults or children with epilepsy. Topiramate is indicated in adults for the prophylaxis of migraine headache.

#### *film-coated tablets*

60 x 25 mg

60 x 50 mg

60 x 100 mg

## URUTAL

### BETAHISTINE

Antivertigo-hemokinetic and otolitic.

#### *tablets*

100 x 8 mg

60 x 16 mg

20 x 24 mg

## VAIRA

### OLANZAPINE

Antipsychotic used to treat the symptoms of psychotic conditions such as schizophrenia and bipolar disorder.

#### *film-coated tablets*

28 x 5 mg

28 x 10 mg

## VAIRA-V

### OLANZAPINE

Antipsychotic used to treat the symptoms of psychotic conditions such as schizophrenia and bipolar disorder.

#### *orodispersible tablets*

28 x 5 mg

28 x 10 mg



## ZARACET

TRAMADOL / ACETAMINOPHEN (PARACETAMOL)

Combination of opioidic analgesic and analgoantipiretic for symptomatic treatment of moderate and severe pain.

*film-coated tablets*

- 10 x (37.5 mg + 325 mg)
- 20 x (37.5 mg + 325 mg)
- 30 x (37.5 mg + 325 mg)
- 100 x (37.5 mg + 325 mg)
- 50 x (75 mg + 650 mg)
- 30 x (75 mg + 650 mg)

P

## ANTIPARASITIC PRODUCTS



## MEDAZOL

METRONIDAZOLE

Antiinfective, antiprotozoal used for treatment of anaerobic infections, protozoal diseases and for prophylaxis in surgical and gynecological procedures.

*tablets*

- 20 x 250 mg
- 10 x 400 mg
- 20 x 400 mg

## RESPIRATORY SYSTEM

## ALERDIN

## DESLORATADINE

Desloratadine is non-sedating long-acting histamine antagonist with selective peripheral H<sub>1</sub>-receptor antagonist activity.

*orodispersible tablets*

10 x 5 mg

30 x 5 mg

*oral solution*

60 ml (0.5 mg / ml)

120 ml (0.5 mg / ml)

## BELODIN / BELODIN A

## LORATADINE

Antihistaminic for systematic use.

Loratadine is used in the treatment of allergy symptoms, such as:

- allergic rhinitis (seasonal and perennial)
- chronic urticaria and other allergic skin disturbances, including neurodermatitis.

*tablets (BELODIN)*

30 x 10 mg

*tablets (BELODIN A)*

7 x 10 mg

14 x 10 mg

## MONLAST

## MONTELUKAST

A leukotriene receptor antagonist (LTRA) used for the maintenance treatment of asthma and to relieve symptoms of seasonal allergies. Montelukast blocks the action of leukotriene D<sub>4</sub> on the cysteinyl leukotriene receptor CysLT<sub>1</sub> in the lungs and bronchial tubes. This reduces the bronchoconstriction otherwise caused by the leukotriene, and results in less inflammation. It is not useful for the treatment of acute asthma attacks.

*chewable tablets*

28 x 5 mg

*film-coated tablets*

28 x 10 mg

## NAXIL

## ACETYLCISTEINE

Mucolytic adjuvant in the therapy of respiratory disorders associated with thick, viscous, mucus hypersecretion. For application in adults and children over the age of 2 years.

*effervescent tablets*

20 x 200 mg

## NAXIL FORTE

### ACETYLCISTEINE

Mucolytic adjuvant in the therapy of respiratory disorders associated with thick, viscous, mucus hypersecretion. For application in adults.

*effervescent tablets*

10 x 600 mg      20 x 600 mg

## RINIL 0.5 mg / ml

### XYLOMETAZOLINE

Application to the nasal passages for children at the age of 2 – 10 years. For the symptomatic relief of nasal congestion, perennial and allergic rhinitis, sinusitis.

*nasal spray, solution*

10 ml

## RINIL 1 mg / ml

### XYLOMETAZOLINE

Application to the nasal passages for adults and children at the age  $\geq 10$  years. For the symptomatic relief of nasal congestion, perennial and allergic rhinitis, sinusitis.

*nasal spray, solution*

10 ml

# FOOD FOR SPECIAL MEDICAL PURPOSES

## NUTRIXA

Only for use under medical supervision

Complete and balanced nutrition. For the dietary management of patients with malnutrition, or at risk of developing malnutrition. High calorie, high protein liquid food with added fiber for a complete, balanced diet. For oral use.

200 ml of high protein liquid food corresponds to 250 kcal.

1 ml of high protein liquid food corresponds to 1.25 kcal.

200 ml of high protein liquid food contain 16 g protein.

Flavour vanilla / Flavour chocolate / Flavour strawberry / Flavour coffee / Flavour peach

200 ml

## NUTRIBEL COMPLEX

For dietary nutrition for patients with malnutrition or the risk of their occurrence, in particular oncological patients, patients with cachexia and chronic catabolic diseases. The preparation is not intended for use in children, except for the recommendation of a physician or other qualified health care provider employees. Caution should be exercised in children aged 3 to 6 years. Not suitable for people with galactosemia, or for a diet without fibers.

High-calorie, high-protein, nutritionally complete liquid-added food EPA , Fiber and Medium Triglycerides. For oral administration. The preparation is provided only with medical supervision.

Flavour chocolate / Flavour cappuccino / Flavour orange-biscuit

300 kcal / 200 ml

20 g protein / 200 ml

1,1 g EPA / 200 ml

## NUTRIBEL DIABET

For the dietary nutrition of malnourished patients (malnutrition associated with the disease) with diabetes or hyperglycemia, or when, for medical reasons, it is not possible to meet the nutritional needs with a normal diet. High-protein, nutritionally complete liquid food with a high content of monounsaturated fatty acids and added fibers. With sugar (fructose) and sweeteners. It contains slowly digestible complex carbohydrates for better regulation of blood glucose levels. Gluten and lactose free. For oral use.

High-calorie, high-protein, nutritionally complete liquid-added food EPA , Fiber and Medium Triglycerides. For oral administration. The preparation is provided only with medical supervision.

Flavour chocolate / Flavour vanilla / Flavour strawberry

200 kcal (844 kJ) / 200 ml

10 g protein / 200 ml

4 g fiber / 200 ml

# Herbal medicines, food supplements, cosmetics and medical devices

## HERBAL MEDICINES

### GINKGO FORTE 120 mg

*Ginkgo bilobae folii extractum siccum*

GINKGO FORTE 120 mg film-coated tablets are intended to improve the condition of patients with impairment of cognitive functions (related to aging) and improvement of the quality of life in patients with mild dementia.

One film-coated tablet contains 120 mg of extract (as dry extract, refined and quantified) from *Ginkgo biloba* L., folium (gingko leaf) (35-67:1), which corresponds to 26.4 – 32.4 mg of flavonoids expressed as flavone glycosides, 3.36 – 4.08 mg of ginkgolide A, B and C and 3.12 – 3.84 mg of bilobalide. Extraction solvent: acetone 65% V/V.

**film-coated tablets**

30 x 120 mg

### HEDERAN syrup

*Hederae helix folii extractum siccum*

Hederan syrup is used as an expectorant in adults and children at the age of 2 and over.

1 ml of syrup contains 3.675 mg of Ivy dry extract (*Hedera helix* L., folium).

**125 ml syrup**

**200 ml syrup**

### SILYMARIN

*Silybi mariani fructi extractum siccum*

SILYMARIN hard capsules are used as supportive therapy in adults to alleviate liver function disorders related to the following diseases and conditions:

- chronic hepatitis B and C
- alcoholic liver disease with cessation of intake of alcoholic beverages
- liver cirrhosis.

One hard capsule contains 168.35 – 205.76 mg of extract (as a dry extract, refined and standardized) from *Silybum marianum* L. Gaertner, fructus (fruit of silymarin) (22–27 : 1) which corresponds to 100 mg of silymarin expressed as silybinin.

Extraction solvent: acetone 95% V/V.

**30 / 60 hard capsules**



## SILYMARIN FORTE

*Silybi mariani fructi extractum siccum*

SILYMARIN FORTE film-coated tablets are used as supportive therapy in adults to alleviate liver function disorders related to the following diseases and conditions:

- chronic hepatitis B and C
- alcoholic liver disease with cessation of intake of alcoholic beverages
- liver cirrhosis.

One film-coated tablet contains 336.70 - 411.52 mg of extract (as dry extract, refined and standardized) from *Silybum marianum* L. Gaertner, fructus (22 - 27:1) which corresponds to 200 mg of silymarin expressed as silibinin .

Extraction solvent: 95% acetone V/V.

**30 film-coated tablets**

## FOOD SUPPLEMENTS

### BELMIRAN DAN (BELMIRAN DAY)

*Passiflora incarnata*

BELMIRAN DAN tablets contain passion flower, which contributes to relaxation of the body.

1 film-coated tablet contains: 100 mg of passion flower dry extract (*Passiflorae incarnatae*).

**20 film-coated tablets**

### BELMIRAN SAN (BELMIRAN NIGHT)

*Valeriana officinalis, Humulus lupulus, Passiflora incarnata*

BELMIRAN SAN tablets help shorten the time needed to initiate and maintain sleep.

1 film-coated tablet contains: 125 mg valerian root dry extract (*Valeriana officinalis*), 30 mg hops strobile dry extract (*Humulus lupulus*) and 80 mg passion flower dry extract (*Passiflora incarnata*).

**20 film-coated tablets**

### FERSAN FORTE tablets

*Carobonyl iron, Vitamin C, Folic acid*

FERSAN FORTE tablets contain iron, vitamin C and folic acid. Iron contributes to normal red blood cell and hemoglobin formation and normal cognitive development of children. Folic acid contributes to normal blood formation and cell division. Vitamin C increases iron absorption.

1 tablet contains: 30 mg (214% RDA) carbonyl iron, 180 mg (225% RDA) vitamin C and 600 µg (300% RDA) folic acid.

**30 tablets**



## MAXI OMEGA-3

### *OMEGA-3 fatty acids*

MAXI OMEGA-3 capsules contain EPA / DHA which contribute to the normal function of the heart.

1 capsule contains 1000 mg of purified and standardized fish oil with a minimum of 30% of omega-3 fatty acids (EPA / DHA), including 18% of eicosapentaenoic acid (EPA) and 12% of docosahexaenoic acid (DHA).

*100 capsules*

## COSMETICS



## BELODEX

### *DEXPANTHENOL 5%*

BELODEX foam contains dexpantenol (provitamin B5) and is used to protect and regenerate dry, sensitive and irritated skin.

It restores and stabilizes the skin's natural protective function and thus prevents water loss, keeping the skin moist and supple. BELODEX foam can be used in conditions of dryness and redness of the skin due to exposure to external influences such as sunlight, heat and cold. In these cases, the application of the foam will bring relief by restoring the elasticity and softness of the skin. The special formulation in the form of foam is extremely gentle and easily spreadable.

BELODEX foam is intended for use in infants, children and adults.

*150 ml*



## EUSIN oil

*Eusyn globulus oil, Mentha piperita oil, Melaleuca leucadendra oil, Menthol, Juniperus communis oil, Methyl salicylate, Eugenia caryophyllus oil*

EUSIN oil is used in baths. EUSIN oil contains mixture of essential oils: eucalyptus oil (*Eucalyptus globulus*), peppermint oil (*Mentha piperita*), cajuput oil (*Melaleuca leucadendra*), juniper oil (*Juniperus communis*) and clove oil (*Eugenia caryophyllus*). It contains also: Methyl salicylate, Menthol, limonene and eugenol.

*drops (10 ml)*



## BELCURA cutaneous spray, emulsion

### *MicroSilver*

Belcura is the Spray Emulsion containing MicroSilver for skin care, protection and regeneration. High-purity MicroSilver provides anti-irritation and anti-bacterial action. Active agents like Ceramides III strengthen the lipid barrier. Glycerin and Lactate hydrate the skin. Hydrophobic lipids form a protective film and deeply nourish the skin without residue. The skin is able to recover and quickly restore its natural barrier.

Ingredients: aqua, butane, caprylic/capric triglyceride, isopentane, cetearyl ethylhexanoate, glycerin, polyglyceryl-3 diisostearate, polyglyceryl-2-dipolyhydroxystearate, dicaprylyl ether, sodium, lactate, magnesium sulfate, silver, ceramide 3, cactic acid, ethylhexylglycerin.

*cutaneous spray (125 ml)*

## EFILIPT BRO cream

*Eucalyptus globulus oil, Melaleuca leucadendron cajaputi oil, Myrtus communis oil*

A warming agent of vegetable origin with expectorant effect. Active essential oils, which are part of the drug, provide warming, bronchodilator, expectorant effect.

Active components: eucalyptus essential oil (Aetheroleum Eucalypti) - 2.0%; cayeputa essential oil (Aetheroleum Melaleuca cajuputi) - 2.0%; essential oil of myrtle (Aetheroleum Myrtus communis) - 2.0%.

Auxiliary components: purified water, PEG-20 methyl glucose sesquistearate, benzyl alcohol, carbomer, sodium hydroxide.

*cream (20 g, 40 g)*

## BELOBAZA plus cream

*with Zinc Oxide and Bisabolol*

BELOBAZA plus is used for regeneration and purification of sensitive and irritated skin. It is also effective for relieving sensations of itching and scratching. Zinc oxide and bisabolol have protective, regenerative and soothing properties are also suitable for keeping the skin in good condition. BELOBAZA plus also contains the emollient white petrolatum which prevents moisture loss while restoring suppleness and flexibility.

Ingredients: aqua, propylene glycol, petrolatum white, cetearyl alcohol, zinc oxide, ceteareth-20, xanthan gum, benzyl alcohol, citric acid monohydrate, phenoxyethanol, bisabolol.

*cream (40 g, 100 g, 400 g)*

## BELOBAZA light cream

*with Zinc gluconate and Bisabolol*

BELOBAZA light cream is used for regeneration of sensitive and irritated skin especially if the skin has undergone various treatments that may affect the natural lipid barrier. Zinc gluconate and bisabolol have protective and soothing properties are also suitable for keeping the skin in good condition while reinforcing the skin's natural barrier. It is also suitable for use in adolescents with blemish prone oily skin because it helps to reduce the appearance of skin pores while purifying the skin. It also mattifies the skin and reduces the appearance of blemishes and redness.

Ingredients: aqua, cyclomethicone, squalene, zinc gluconate, propylene glycol, ceteareth-20, xanthan gum, benzyl alcohol, bisabolol, sodium hydroxide.

*cream (50 g, 60 g, 100 g)*

## BELOSALIC SALIC

*with 3% salicylic acid anti-dandruff shampoo*

BELOSALIC SALIC shampoo contains an effective formulation that is designed to reduce, eliminate and prevent the reformation of dandruff, while providing a soothing and regenerating effect on the skin leaving the hair and scalp clean and smooth with a sensation of freshness.

The formulation is based on 3% salicylic acid and piroctone olamine. Salicylic acid is effective at cleaning the upper layers of the skin on the scalp from dandruff and excessive sebum thus eliminating visible flakes. It also softens the skin and helps to maintain the natural acidity of the skin.

Piroctone olamine soothes the scalp and reduces flaking while preventing the formation of new dandruff. It helps the scalp stay clean and without irritation.

Panthenol and chamomile flower extract act soothing and restore the natural skin barrier properties thus providing gentle care for the hair and scalp.

*Packaging size 100 ml / 200 ml*

# MEDICAL DEVICES

## BELOBAZA cream

### SOFT PARAFIN AND FAT PRODUCTS

BELOBAZA regenerates and helps maintaining physiological balance of the skin – it protects, moisturizes, soothes and calms skin, as well as regenerates. BELOBAZA contains no parabens, synthetic dyes and parfum and it is therefore suitable for care and protection of sensitive skin, including sensitive skin of children. Due to adjusted pH value, BELOBAZA helps maintaining natural acid pH of the skin.

Ingredients: aqua, petrolatum, cetearyl alcohol, paraffinum liquidum, ceteareth-20, benzyl alcohol, sodium phosphate, phosphoric acid, sodium hydroxide.

*cream*

100 g      400 g

# AUXILIARY SUBSTANCES

## BELOBAZA BELUPO

### SOFT PARAFIN AND FAT PRODUCTS

Ambiphilic base of carefully balanced ingredients for skin treatment, protection and care, and for preparation of magistral medicines (can be combined with almost any dermatotherapeutic).

*cream*

2 kg      4 kg

# Contacts

***Belupo is strategically oriented towards increasing exports and developing new drugs that meet the needs of demanding foreign markets. Our organization and work ethic is flexibly adjusted to contemporary business management practice, both at home and on foreign markets.***



**Belupo, lijekovi i kozmetika, d.d.**

Danica 5,  
48 000 KOPRIVNICA

T: +385 48 659 000

T: +385 (1) 24 81 222

E- mail: [belupo@belupo.hr](mailto:belupo@belupo.hr)

[www.belupo.hr](http://www.belupo.hr)

[www.belupo.com](http://www.belupo.com)

[www.zdravobudi.hr](http://www.zdravobudi.hr)

[www.neofen.hr](http://www.neofen.hr)

[www.lupocet.hr](http://www.lupocet.hr)



# Croatia

## BOARD OF DIRECTORS OFFICES

Ulica Danica 5  
48 000 Koprivnica  
Phone: +385 (48) 652 200  
E- mail: belupo@belupo.hr

I. Savica 36, P.P. 62  
10 000 Zagreb  
Phone: +385 (1) 24 81 222  
E- mail: belupo@belupo.hr

## MARKET OF THE REPUBLIC OF CROATIA

### Marketing Pharmaceuticals

I. Savica 36, P.P. 62  
10 000 Zagreb  
Phone: +385 (1) 24 81 222  
E- mail: belupo@belupo.hr

### Sales

Josipa Vargovića 4/3  
48000 Koprivnica  
Phone: +385 (48) 659 032  
+385 (48) 659 033  
E- mail: belupo@belupo.hr

I. Savica 36, P.P. 62  
10 000 Zagreb  
Phone: +385 (1) 24 81 222  
E- mail: belupo@belupo.hr

### Pharmacies Deltis Pharm

OFFICIES  
Josipa Vargovića 4/IV  
48000 Koprivnica  
Phone: +385 (48) 659 015  
E- mail: belupo@belupo.hr

### Purchase

Ulica Danica 5, 48000 Koprivnica  
Phone: +385 (48) 659 005  
E- mail: belupo@belupo.hr

## INTERNATIONAL MARKETS

Ulica Danica 5  
48000 Koprivnica  
Phone: +385 (48) 659 002  
E- mail: belupo@belupo.hr

## PRODUCTION

Ulica Danica 5,  
48000 Koprivnica  
Phone: +385 (48) 652 200  
E- mail: belupo@belupo.hr

## RESEARCH AND DEVELOPMENT

Ulica Danica 5, 48000 Koprivnica  
Phone: +385 (48) 652 523  
E- mail: belupo@belupo.hr

## BUSINESS DEVELOPMENT AND REGULATORY AFFAIRS

I. Savica 36, P.P. 62, 10000 Zagreb  
Phone: +385 (1) 24 81 266  
E- mail: belupo@belupo.hr

## HUMAN RESOURCES

Ulica Danica 5, 48000 Koprivnica  
Phone: +385 (48) 652 534,652 204  
E- mail: kadrovi@belupo.hr  
E- mail: ljudski.potencijali@belupo.hr

## LEGAL AND GENERAL AFFAIRS

Ulica Danica 5, 48000 Koprivnica  
Phone: +385 (48) 652 769  
E- mail: belupo@belupo.hr

## PUBLIC RELATIONS

I. Savica 36, P.P. 62, 10 000 Zagreb  
Phone: +385 (1) 24 81 292  
E- mail: public.relations@belupo.hr

# Companies

## SLOVENIA

### **BELUPO d.o.o.**

Dvoržakova 6 | 1000 Ljubljana  
Phone: +386 1 300 9511  
E-mail: info@belupo.si

## SLOVAKIA

### **BELUPO S.R.O.**

Cukrová 14 / 210 | 811 08 Bratislava  
Phone: +421 2 59 32 43 30  
E-mail: belupo@belupo.sk

## NORTH MACEDONIA

### **BELUPO D.O.O.E.L. SKOPJE**

Ul. 3-ta Makedonska brigada 68 |  
1000 Skopje  
Phone: +389 22 403 783  
E-mail: belupo.mk@belupo.com.mk

## BOSNIA AND HERZEGOVINA

### **FARMAVITA d.o.o.**

Igmanska 5a  
71 320 Vogošća  
Phone: +387 33 476 320  
E-mail: farmavita@farmavita.ba

# Representative offices

## BOSNIA AND HERZEGOVINA

Hasana Merdžanovića bb  
71 320 Vogošća  
Phone: +387 33 426 615; 426 616  
E-mail: belupo@bih.net.ba

## CZECH REPUBLIC

BELUPO léky a kosmetika, s.r.o.  
Průmyslová 1306/7, Hostivař,  
102 00 Praha 10  
Phone: +420 2 74 77 04 14  
E-mail: belupo@belupo.cz

## KOSOVO

BELUPO-PERFAQESIA PRISHTINE  
Mati 1, Rr.B, H-A, Nr. 3, Residio 2  
10000 Prishtine, KOSOVO

## RUSSIAN FEDERATION

Lomonosovsky pr. 38,  
app. 68 (office-manager) 71-72, 124  
119330 Moscow  
Phone: +7 495 933 7212 | 7213 | 7214  
E-mail: belupo@belupo.su

## SERBIA

Mosorska 1 | 11 000 Beograd  
Phone: +381 11 30 98 871, 30 98 872  
E-mail: office@belupo.co.rs

## UKRAINE

18 Chigorina str., office 302  
01042 Kiev  
Phone: +38 067 465 17 31  
E-mail: office@belupo.com.ua

## KAZAHSTAN

Timityazev str. 42, off. 328  
050000 Almaty  
Republik of Kazakhstan  
Phone: + 7 727 245 88 42  
E-mail: belupo@belupo.kz