

# ABOUT US



Pharmaceuticals & Cosmetics  
Koprivnica, Croatia  
[www.belupo.com](http://www.belupo.com)  
e-mail: [belupo@belupo.hr](mailto:belupo@belupo.hr)

Belupo, the second-largest pharmaceutical producer in Croatia, is the country's leading seller of both cardiovascular and dermatological drugs.

Belupo was founded in 1971 and its first production plant in Ludbreg became operational in 1972. Rapid development and product range expansion brought about the construction of another plant in Koprivnica in 1980, where the company is still headquartered today. Continuous investment in production modernization and employee education, implementation of scientific methods and assurance of product high quality standards, as well as dedicated team work resulted in the development of new products, market share increase and export sales volume growth, positioning Belupo among the leaders.

In 1999 Belupo opened new solid pharmaceuticals facility which is the largest and the most important production site in the company's 20,000 m<sup>2</sup> pharmaceutical complex. In 2004 a new facility for the production of semi-solid and liquid pharmaceuticals was opened. Belupo opened a new Quality Control and QA facilities in 2005. The newest of Belupo's investments, the construction of the new factory for semi-liquid, liquid, solid and sterile product forms will be completed in 2017.

The new plants meet top standards of modern pharmaceutical industry, including the Good Manufacturing Practice Standards, the U.S. Food and Drug Administration requirements, The European Pharmaceutical Inspection Convention regulations, and the requirements and standards set in the Croatian Drug Law.

At Belupo we develop, improve and produce human pharmaceuticals, traditional herbal preparations and herbal medicinal products and food supplement. Numerous and diverse forms of cooperation include reputable foreign business partners: Mylan, Dermatologie, Pharmaten, Medis, Lupin, Torrentconfirm and guarantee our dedication to latest developments in the field of pharmaceuticals.

According to the anatomical-therapeutic-chemical (ATC) classification, Belupo's pharmaceuticals fall under eleven groups. Belupo's product portfolio is focused on the following: cardiovascular system medicines, musculo-skeletal system medicines, nervous system medicines, dermatological and general anti-infective medicines for systemic use.

# CONTACTS

## CROATIA

### ● BOARD OF DIRECTORS OFFICES

I. Savica 36 | P.P. 62  
10 000 Zagreb  
Phone: +385 1 24 81 222  
Fax: +385 1 23 71 411  
E-mail: belupo@belupo.hr

Josipa Vargovića 4/3  
48 000 Koprivnica  
Phone: +385 48 659 011  
Fax: +385 48 624 992  
E-mail: belupo@belupo.hr

Opatička 5  
48000 Koprivnica  
Phone: +385 48 659 000  
Fax: +385 48 624 271  
E-mail: belupo@belupo.hr

### ● PUBLIC RELATION

I. Savica 36 | P.P. 62, 10 000 Zagreb  
Phone: +385 1 24 81 201  
Fax: +385 1 23 71 411  
E-mail: public.relations@belupo.hr

### ● BUSINESS DEVELOPMENT AND REGULATORY AFFAIRS

I. Savica 36 | P.P. 62, 10 000 Zagreb  
Phone: +385 1 24 81 266  
Fax: +385 1 24 81 290  
E-mail: belupo@belupo.hr

### ● MARKETING PHARMACEUTICALS

I. Savica 36 | P.P. 62, 10 000 Zagreb  
Tel +385 1 24 81 210  
Fax +385 1 23 71 370  
E-mail: belupo@belupo.hr

### ● MARKETING SELF-MEDICATION PRODUCTS

I. Savica 36 | P.P. 62, 10 000 Zagreb  
Tel +385 1 24 81 203  
Fax +385 1 23 71 411  
E-mail: bezreceptni@belupo.hr

### ● PROMOTION

I. Savica 36 | P.P. 62, 10 000 Zagreb  
Tel +385 1 24 81 247  
Fax +385 1 23 71 396  
E-mail: promocija@belupo.hr

### ● INTERNATIONAL MARKETS

I. Savica 36 | P.P. 62, 10 000 Zagreb  
Phone: +385 1 24 81 209  
Fax: +385 1 23 71 399  
E-mail: belupo@belupo.hr

### ● SALES & LOGISTIC

Josipa Vargovića 4/3  
48000 Koprivnica  
Phone: +385 48 659 030; +385 48 659 032  
Fax: +385 48 624 065  
E-mail: belupo@belupo.hr

I. Savica 36 | P.P. 62  
10000 Zagreb  
Phone: +385 1 24 81 211  
Fax: +385 1 23 71 411  
E-mail: belupo@belupo.hr

### ● QUALITY CONTROL

Danica 5, 48000 Koprivnica  
Phone: +385 48 652 209  
Fax: +385 48 647 085  
E-mail: belupo@belupo.hr

### ● INFORMATION TECHNOLOGY

Danica 5, 48000 Koprivnica  
Phone: +385 48 652 600  
Fax: +385 48 652 214  
E-mail: belupo@belupo.hr

### ● QUALITY MANAGEMENT

Danica 5, 48000 Koprivnica  
Phone: +385 48 652 891  
Fax: +385 48 652 894  
E-mail: belupo@belupo.hr

### ● LEGAL AND GENERAL AFFAIRS

Danica 5, 48000 Koprivnica  
Phone: +385 48 652 212  
Fax: +385 48 652 531  
E-mail: belupo@belupo.hr

# CONTACTS

## CROATIA

### ● PRODUCTION

Danica 5, 48000 Koprivnica  
Phone: +385 48 652 200  
Fax +385 48 647 087  
E-mail: belupo@belupo.hr

### ● PURCHASE

Josipa Vargovića 4, 48000 Koprivnica  
Phone: +385 48 659 005  
Fax: +385 48 624 996  
E-mail: belupo@belupo.hr

### ● CONTROLLING

Danica 5  
48000 Koprivnica  
Phone: +385 48 652 244  
Fax: +385 48 652 544  
E-mail: belupo@belupo.hr

### ● TREASURY AND ACCOUNTING

|  |   |
|--|---|
| Opatička 5<br>48000 Koprivnica<br>Phone: +385 48 659 024<br>Fax: +385 48 622 879<br>E-mail: belupo@belupo.hr | Josipa Vargovića 4/1<br>48000 Koprivnica<br>Phone: +385 48 659181<br>Fax: +385 48 659 194<br>E-mail: belupo@belupo.hr |
|--|---|

# CONTACTS

## COMPANIES & REPRESENTATIVE OFFICES ABROAD

Belupo is strategically oriented towards increasing exports and developing new drugs that meet the needs of demanding foreign markets. Our organization and work ethic is flexibly adjusted to contemporary business management practice, both at home and on foreign markets.

Our foreign market operations are supported by a network of representative offices Sarajevo, Skopje, Belgrade, Prague, Bratislava, Bucharest and Moscow and companies in Ljubljana (Slovenia) and Bratislava (Slovakia).

## COMPANIES

### ● SLOVENIA

BELUPO d.o.o.  
Dvoržakova 6 | 1000 Ljubljana  
Phone: +386 1 300 9510  
Fax: +386 1 432 6311  
E-mail: info@belupo.si

### ● SLOVAKIA

BELUPO S.R.O.  
Cukrová 14 / 210 | 811 08 Bratislava  
Phone: +421 2 59 32 43 30; +421 2 59 32 43 34  
Fax: +421 2 59 32 43 31  
E-mail: pavel\_petrovic@belupo.sk

### ● MACEDONIA

BELUPO DOEL SKOPJE  
Ul. 3-ta Makedonska udarna brigada 68 | 1000 Skopje  
Phone: +389 22 403 783  
Fax: +389 22 403 790  
E-mail: belupo@belupo.com.mk

### ● BOSNIA AND HERZEGOVINA

FARMAVITA d.o.o.  
Igmanska 5a | 71 320 Vogošća  
Phone: +387 33 476 320  
Fax: +387 33 476 321  
E-mail: farmavita@farmavita.ba

# CONTACTS

## REPRESENTATIVE OFFICES

### ● BOSNIA AND HERZEGOVINA

Hasana Merdžanovića bb  
71 320 Vogošća  
Phone: +387 33 426 615; 426 616  
Fax: +387 33 426 617  
E-mail: belupo@bih.net.ba

### ● KOSOVO

BELUPO-PERFAQESIA PRISHTINE  
Mati 1, Rr.B, H-A, Nr. 3, Residio 2  
10000 Prishtine, KOSOVO  
Phone/Fax: + 381 38 70 45 16  
E-mail: belupo.kosova@ifs-ks.net

### ● SERBIA

Mosorska 1 | 11 000 Beograd  
Phone: +381 11 30 98 871, 30 98 872  
Fax: +381 11 24 93 226  
E-mail: belupo@eunet.rs

### ● POLAND

Ul. Śniadeckich 10/20 | 00-656 Warsaw  
Cell: +3848 72 65 86 678  
E-mail: office@belupo.pl

### ● UKRAINE

18 Chigorina str., office 302 | 01042 Kiev  
Phone: +38 044 285 87 20  
Fax: +38 044 285 87 22  
E-mail: office@belupo.com.ua

### ● CZECH REPUBLIC

Černokostelecká 9  
100 00 Prague  
Phone: +420 2 74 77 04 14  
Fax: +420 2 74 77 20 11  
E-mail: belupo@belupo.cz

### ● RUSSIAN FEDERATION

Lomonoskovsky pr. 38, app. 71-72, 124  
119330 Moscow  
Phone: +7 495 933 7212 | 7213 | 7214  
Fax: +7 495 933 7215  
E-mail: belupo@belupo-dd.ru

### ● SLOVAKIA

BELUPO S.R.O.  
Cukrová 14 / 210 | 811 08 Bratislava  
Phone: +421 2 59 32 43 30; 59 32 43 32  
Fax: +421 2 59 32 43 31  
E-mail: pavel\_petrovic@belupo.sk

### ● KAZAHSTAN

Zheltaksan str.87, off. 24,25, 26  
050000 Almaty  
Republik of Kazahstan  
Phone: + 7 771 506 06 18  
Fax: + 7 727 279 35 33  
E-mail: belupo@belupo.kz

# PRODUCT RANGE

## PHARMACEUTICALS



A

## ALIMENTARY TRACT AND METABOLISM

### BELFORMIN<sup>®</sup>

METFORMINE

Antidiabetic drug, a biguanide antihyperglycemic agent used for treating non-insulin-dependent diabetes mellitus (NIDDM). Metformin improves glycemic control by decreasing hepatic glucose production, decreasing glucose absorption and increasing insulin-mediated glucose uptake.

*film-coated tablets*

|              |              |               |
|--------------|--------------|---------------|
| 100 x 500 mg | 30 x 850 mg  | 30 x 1000 mg  |
|              | 120 x 850 mg | 120 x 1000 mg |

### BELOSEPT<sup>®</sup>

HEXETIDINE

Oral antiseptic, a broad-spectrum antibacterial and antimycotic agent for rinsing the mouth.

*solution*

200 ml (1 mg / ml)

## ● DIAPIRID®

### GLIMEPIRIDE

Antidiabetic drug, the first III generation sulphonylurea used for treating non-insulin-dependent diabetes mellitus (NIDDM) with long duration of action.

The mechanism of action of glimepiride in lowering blood glucose appears to be dependent on stimulating the release of insulin from functioning pancreatic beta cells, and increasing sensitivity of peripheral tissues to insulin.

#### *tablets*

30 x 1 mg          30 x 2 mg          30 x 3 mg

## ● GASTROBEL®

### RANITIDINE

Specific rapidly acting histamine H<sub>2</sub>-receptor antagonist. Used for symptomatic relief of heartburn, indigestion, acid indigestion and hyperacidity.

#### *effervescent tablets*

10 x 150 mg

## ● GLIKA®

### GLICLAZIDE

Hypoglycemic sulphonylurea oral antidiabetic active substance. Reduces blood glucose levels by stimulating insulin secretion from β-cells of the islands of Langerhans in pancreas.

#### *prolonged-release tablets*

60 x 30 mg

## ● LORDIAR®

### LOPERAMIDE

Synthetic antidiarrheal for oral use.

#### *capsules*

20 x 2 mg

## ● PORTALAK® efekt

### LACTULOSE

Osmotic laxative, used in occasional or long-term constipation.

#### *syrup*

250 ml (66.7 g / 100 ml)

## ● PORTALAK®

### LACTULOSE

Osmotic laxative, used in occasional or long-term constipation.

#### *syrup*

500 ml (66.7 g / 100 ml)

## POTASSIUM CHLORIDE BELUPO (KALIJEV KLORID BELUPO)

### POTASSIUM CHLORIDE

Oral replacement of electrolyte and fluid loss due to various reasons.

#### *soluble tablets*

30 x 500 mg

## REGLINID<sup>®</sup>

### REPAGLINIDE

Antidiabetic drug, an oral blood glucose-lowering drug of the meglitinide class, used in the management of non-insulin-dependent diabetes mellitus (NIDDM). Repaglinide lowers blood glucose levels by stimulating the release of insulin from the pancreas.

#### *tablets*

90 x 0.5 mg

90 x 1 mg

90 x 2 mg

## ROJAZOL<sup>®</sup>

### MICONAZOLE NITRATE

Topical broad-spectrum antimycotic. Combined fungistatic and fungicidal action, bacteriostatic and bactericidal action on gram-positive cocci and bacilli that often follow fungous infections

#### *oral gel*

40 g 2% (20 mg / g)

## RUDAKOL<sup>®</sup>

### MEBEVERINE

Selective musculotropic spasmolytic without anticholinergic side effects.

#### *coated tablets*

50 x 135 mg

## TARGET<sup>®</sup> PLUS

### OMEPRAZOLE

Proton pumps inhibitor, effective in the short-term treatment of duodenal, gastric and esophageal ulcers and may offer a useful alternative to conventional therapy. In combination with antimicrobials in dual and triple drug regimens, omeprazole therapy can achieve high cure rates in the treatment of *Helicobacter pylori* associated with peptic ulcer disease. The drug is recommended for Zollinger-Ellison syndrome.

#### *capsules*

14 x 20 mg

28 x 20 mg

## ZOLTEX<sup>®</sup>

### PANTOPRAZOLE

Proton pumps inhibitor, effective in the short-term treatment of duodenal, gastric and esophageal ulcers and may offer a useful alternative to conventional therapy. In combination with antimicrobials in dual and triple drug regimens, omeprazole therapy can achieve high cure rate in the treatment of *Helicobacter pylori* associated with peptic ulcer disease. The drug is recommended for Zollinger-Ellison syndrome.

#### *gastro-resistant tablets*

28 x 20 mg

14 x 40 mg

56 x 20 mg

28 x 40 mg

#### *powder for solution for injection*

40 mg / 10 ml

# B

## BLOOD AND BLOOD FORMING ORGANS

### ● ASKA<sup>®</sup> pro

#### ACETYLSALICYLIC ACID

Acetylsalicylic acid inhibits the adhesion and dumping of blood platelets (thrombocytes) and in this way prevents blood clots (thrombi) to develop (inhibition of platelet aggregation).

*gastro-resistant tablets*

30 x 100 mg

### ● HEPARIN BELUPO

#### HEPARIN

Endogenous anticoagulant for intravenous infusion or intravenous injection.

*solution for injections*

10 x 5 ml (25,000 IU / 5 ml)

### ● KLOPIDEX<sup>®</sup>

#### CLOPIDOGREL

Clopidogrel is an oral antiplatelet agent (thienopyridine class) which inhibits blood clotting and is indicated for prevention of vascular ischaemic events in patients with symptomatic atherosclerosis, acute coronary syndrome without ST-segment elevation (NSTEMI), ST elevation MI (STEMI). It is also used, along with aspirin, for the prevention of thrombosis after placement of intracoronary stent or as an alternative antiplatelet drug for patients who are intolerant to aspirin.

*film-coated tablets*

30 x 75 mg



## CARDIOVASCULAR SYSTEM

## ● AMIODARON BELUPO

## AMIODARONE

Antiarrhythmic drug, class III.

*tablets*

60 x 200 mg

● AMONEX<sup>®</sup>

## AMLODIPINE

Third generation of the dihydropyridine derivative. Calcium antagonist with predominantly vascular effect, indicated for the treatment of essential hypertension, chronic stable angina pectoris and Prinzmetal angina.

*tablets*

30 x 5 mg

60 x 5 mg

90 x 5 mg

30 x 10 mg

60 x 10 mg

90 x 10 mg

● BISOBEL<sup>®</sup>

## BISOPROLOL

Cardioselective  $\beta_1$ -adrenergic receptor blocker, indicated for treatment of essential hypertension, coronary heart disease and congestive heart failure.

*tablets*

30 x 2.5 mg

30 x 5 mg

60 x 2.5 mg

60 x 5 mg

30 x 10 mg

● CARVELOL<sup>®</sup>

## CARVEDILOL

$\alpha$ - and  $\beta$ -adrenergic receptor blocker, indicated for the treatment of essential hypertension, coronary heart disease and symptomatic congestive heart failure.

*tablets*

30 x 3.125 mg

30 x 6.25 mg

30 x 12.5 mg

30 x 25 mg

## DOXA<sup>®</sup>

### DOXAZOSIN

Alpha-adrenergic blockers. It causes the blood vessels (veins and arteries) to relax and expand, improving blood flow. Doxazosin also relaxes muscles in the prostate and bladder neck, making it easier to urinate.

**tablets**

20 x 2 mg            20 x 4 mg

## FURSEMID

### FUROSEMIDE

Potent loop diuretic. Inhibits sodium and chloride reabsorption at the loop of Henle.

**tablets**

20 x 40 mg

**solution for injections**

5 x 2 ml (20 mg / 2 ml)

## FURSEMID FORTE

### FUROSEMIDE

Potent loop diuretic. Inhibits sodium and chloride reabsorption at the loop of Henle.

**tablets**

20 x 500 mg

## INDAMID<sup>®</sup> SR

### INDAPAMIDE

Diuretic and antihypertensive. Pharmacologically related to thiazide diuretics with antihypertensive action.

**prolonged-release tablets**

30 x 1.5 mg

## IRUMED<sup>®</sup>

### LISINOPRIL

ACE inhibitor, used in hypertension, heart failure and for short-term treatment hemodynamically stable patients after acute myocardial infarction. Slows down the development of diabetic nephropathy and retinopathy.

**tablets**

30 x 2.5 mg

30 x 5 mg

30 x 10 mg

30 x 20 mg

60 x 2.5 mg

60 x 5 mg

60 x 10 mg

60 x 20 mg

## IRUZID<sup>®</sup>

### LISINOPRIL AND HYDROCHLOROTHIAZIDE

Combination of ACE inhibitor and diuretic, antihypertensive

**tablets**

30 x (10 + 12.5 mg)

30 x (20 + 12.5 mg)

30 x (20 + 25 mg)

60 x (10 + 12.5 mg)

60 x (20 + 12.5 mg)

60 x (20 + 25 mg)

## ISOCARD<sup>®</sup>

### ISOSORBIDE MONONITRATE

Used in prevention of angina pectoris attacks.

#### *prolonged-release tablets*

50 x 40 mg

50 x 60 mg

## LACIDIPIN BELUPO

### LACIDIPINE

Calcium channel blocker, indicated in adults for the treatment of hypertension, either alone or in combination with other antihypertensive agents.

#### *film-coated tablets*

28 x 4 mg

28 x 6 mg

## MOKSONIDIN BELUPO

### MOKSONIDINE

Antihypertensive centrally acting antiadrenergic agent.

#### *film-coated tablets*

28 x 0.2 mg

28 x 0.4 mg

## NIBEL<sup>®</sup>

### NEBIVOLOL

Antihypertensive and selective  $\beta$ 1-blocker with nitric oxide-potentiating vasodilatory effect, indicated for treatment of essential hypertension and congestive heart failure.

#### *tablets*

14 x 5 mg

28 x 5 mg

## ORMIDOL<sup>®</sup>

### ATENOLOL

Antihypertensive and selective  $\beta$ -blocker

#### *film-coated tablets*

30 x 25 mg

30 x 50 mg

28 x 100 mg

## PINOX<sup>®</sup>

### LERCANIDIPINE

Third generation of the dihydropyridine derivative. Calcium antagonist with predominantly vascular effect, indicated for the treatment of essential hypertension. Lercanidipine works by relaxing and opening the blood vessels allowing the blood to circulate more freely around the body. This lowers the blood pressure and allows the heart to work more efficiently.

#### *film-coated tablets*

28 x 10 mg

56 x 10 mg

## PROTECTA<sup>®</sup>

### SIMVASTATINE

Hypolipemic, an HMG-CoA reductase inhibitor.

*film-coated tablets*

|            |            |            |
|------------|------------|------------|
| 28 x 10 mg | 28 x 20 mg | 28 x 40 mg |
|            | 56 x 20 mg |            |

## RAMED<sup>®</sup>

### RAMIPRIL

Antihypertensive. ACE inhibitor, indicated for the treatment of hypertension, heart failure and post-myocardial infarction. Slows down the development of non-diabetic and diabetic nephropathy.

*tablets*

|              |             |           |            |
|--------------|-------------|-----------|------------|
| 28 x 1.25 mg | 28 x 2.5 mg | 28 x 5 mg | 28 x 10 mg |
|              | 56 x 2.5 mg | 56 x 5 mg |            |

## RAMZID<sup>®</sup>

### RAMIPRIL AND HYDROCHLOROTHIAZIDE

Antihypertensive. Combination of ACE inhibitor and diuretic.

*tablets*

|                      |                  |
|----------------------|------------------|
| 28 x (2.5 + 12.5 mg) | 28 x (5 + 25 mg) |
| 56 x (2.5 + 12.5 mg) | 56 x (5 + 25 mg) |

## ROSIX<sup>®</sup>

### ROSUVASTATIN

Hypolipemic, an inhibitor of HMG-CoA reductase, reduces cholesterol by increasing the number of low-density lipoprotein (LDL) receptors on the cell-surface to enhance uptake and catabolism of LDL, inhibits hepatic synthesis of hepatic very-low-density lipoprotein (VLDL), which reduces the total number of VLDL and LDL particles. The treatment reduces triglycerides (TG) and produces increases in high-density lipoprotein cholesterol (HDL).

*film-coated tablets*

28 x 10 mg  
28 x 20 mg  
28 x 40 mg

## TOMID<sup>®</sup>

### TORASEMIDE

Diuretic and antihypertensive. Torasemide is a pyridine-sulfonyl urea type loop diuretic mainly used in the management of edema associated with congestive heart failure. It is also used at low doses for the management of hypertension.

*tablets*

30 x 5 mg  
30 x 10 mg

## TORVAS<sup>®</sup>

### ATORVASTATINE

Hypolipemic. Indicated as an adjunct to diet for reduction of elevated total cholesterol, LDL-cholesterol, apolipoprotein B, and triglycerides in patients with primary hypercholesterolaemia, heterozygous familial hypercholesterolaemia or combined (mixed) hyperlipidaemia when response to diet and other nonpharmacological measures is inadequate.

#### *film-coated tablets*

|            |            |            |
|------------|------------|------------|
| 30 x 10 mg | 30 x 20 mg | 30 x 40 mg |
| 60 x 10 mg | 60 x 20 mg | 60 x 40 mg |

## VAL<sup>®</sup>

### VALSARTAN

Antihypertensive. Angiotensin II antagonist acting on the AT1 receptor, indicated for the treatment of hypertension, heart failure and post-myocardial infarction.

#### *film-coated tablets*

|            |             |
|------------|-------------|
| 28 x 80 mg | 28 x 160 mg |
| 56 x 80 mg | 56 x 160 mg |

## VAL<sup>®</sup> PLUS

### VALSARTAN AND HYDROCHLOROTHIAZIDE

Antihypertensive. Combination of Angiotensin II antagonist and diuretic.

#### *film-coated tablets*

|                     |                      |
|---------------------|----------------------|
| 28 x (80 + 12.5 mg) | 28 x (160 + 12.5 mg) |
| 56 x (80 + 12.5 mg) | 56 x (160 + 12.5 mg) |

D

## DERMATOLOGICALS

## AFLODERM<sup>®</sup>

### ALCLOMETHASONE DIPROPIONATE

Non-fluorinated corticosteroid dermatological with antiinflammatory, antiallergic and vasoconstrictive action, suitable for use in all age groups.

#### *cream*

|                   |
|-------------------|
| 20 g (0.5 mg / g) |
| 40 g (0.5 mg / g) |

#### *ointment*

|                   |
|-------------------|
| 20 g (0.5 mg / g) |
|-------------------|

## ALOMAX<sup>®</sup>

### MINOXIDIL

Treatment of androgenic alopecia in men and women between 18 and 65 years. Alomax solution works as a vasodilator and therefore improves blood circulation in the scalp.

#### **solution**

60 ml 5% (for men only)

60 ml 2% (for men and women)

## AMOFIN<sup>®</sup>

### AMOROLFINE

Broad spectrum antimycotic for topical use. Used for treatment of onychomycosis without matrix involvement, caused by dermatophytes, yeasts or moulds.

#### **medicated nail lacquer**

2.5 ml (50 mg / ml)

the pack contains 30 cleansing swabs,  
10 spatulas and 30 nail files

5 ml (50 mg / ml)

the pack contains 30 cleansing swabs,  
10 spatulas and 60 nail files

## BELODERM<sup>®</sup>

### BETAMETHASONE DIPROPIONATE

Fluorinated corticosteroid dermatological with antiinflammatory, antipruriginous, antiallergic and vasoconstrictive action, suitable for treatment of chronic and persistent dermatoses.

#### **cream**

15 g (0.5 mg / g)

30 g (0.5 mg / g)

100 g (0.5 mg / g)

#### **ointment**

15 g (0.5 mg / g)

30 g (0.5 mg / g)

#### **skin solution**

50 ml (0.5 mg / g)

## BELOGENT<sup>®</sup>

### BETAMETHASONE DIPROPIONATE AND GENTAMICIN SULPHATE

Fluorinated corticosteroid dermatological combined with a wide-spectrum antibiotic, suitable for treatment of primary and secondary infected dermatoses.

#### **cream**

15 g (0.5 + 1 mg / g)

#### **ointment**

15 g (0.5 + 1 mg / g)

## BELOSALIC<sup>®</sup>

### BETAMETHASONE DIPROPIONATE AND SALICYLIC ACID

Fluorinated corticosteroid dermatological with keratolytic, suitable for treatment of dry, peeling and hyperkeratotic dermatoses.

#### **ointment**

30 g (0.5 + 30 mg / g)

#### **skin solution**

50 ml (0.5 + 20 mg / g)

100 ml (0.5 + 20 mg / g)

## HERPLEX<sup>®</sup>

### ACYCLOVIR

Highly selective antiviral for treatment of infections caused by herpes simplex and varicella-zoster viruses.

#### **cream**

2 g (50 mg / g)

## HERPLEXIM<sup>®</sup>

### ACYCLOVIR

Highly selective antiviral for treatment of infections caused by herpes simplex and varicella-zoster viruses.

*cream*

5 g (50 mg / g)                      10 g (50 mg / g)

## MIROBACT<sup>®</sup>

### MUPIROCIN

Mupirocin is used as a topical treatment for primary and secondary bacterial skin infections.

*ointment*

15 g 2% (20 mg / g)

## ROJAZOL<sup>®</sup>

### MICONAZOLE NITRATE

Topical broad-spectrum antimycotic. Combined fungistatic and fungicidal action, and bacteristatic and bactericidal action on gram-positive cocci and bacilli that often follow fungous infections.

*cream*

30 g 2% (20 mg / g)

## SONA<sup>®</sup>

### ADAPALENE

Topical retinoid suitable for treatment of mild-moderate acne.

*gel*

30 g (1 mg / g)

*cream*

30 g (1 mg / g)

## TERBINAX<sup>®</sup>

### TERBINAFINE

Antimycotic. Terbinafine is used to treat certain types of fungal infection. It works by stopping the growth of fungus.

*tablets*

14 X 250 mg

42 X 250 mg

## GENITO URINARY SYSTEM AND SEX HORMONES

**MEDAZOL<sup>®</sup>**

## METRONIDAZOLE

Antiinfective, antiprotozoal used for treatment of anaerobic infections, protozoal diseases and for prophylaxis in surgical and gynecological procedures.

*vaginal tablets*

10 x 500 mg

**ROJAZOL<sup>®</sup>**

## MICONAZOLE NITRATE

Topical broad-spectrum antimycotic. Combined fungistatic and fungicidal action, bacteriostatic and bactericidal action on gram-positive cocci and bacilli that often follow fungus infections.

*vaginal suppositories*

7 x 200 mg

**TAMOSIN<sup>®</sup>**

## TAMSULOSIN

An antiadrenergic agent. It is used to treat enlargement of the prostate gland in men, a condition called benign prostatic hyperplasia or BPH.

*prolonged-release capsules*

30 x 0.4 mg

**VINER<sup>®</sup>**

## SILDENAFIL

In the presence of sexual stimulation, sildenafil enhances the effect of nitric oxide (NO) by inhibiting phosphodiesterase type 5 (PDE5), which is responsible for degradation of cyclic guanosine monophosphate (cGMP) in the corpus cavernosum.

*film-coated tablets*

4 x 50 mg

4 x 100 mg

**VINER<sup>®</sup> MINT**

## SILDENAFIL

In the presence of sexual stimulation, sildenafil enhances the effect of nitric oxide (NO) by inhibiting phosphodiesterase type 5 (PDE5), which is responsible for degradation of cyclic guanosine monophosphate (cGMP) in the corpus cavernosum.

*chewable tablets*

4 x 50 mg

4 x 100 mg



## GENERAL ANTIINFECTIVES FOR SYSTEMIC USE

## ● AMOKSICILIN BELUPO

## AMOXICILLIN TRIHYDRATE

Semi-synthetic penicillin wide-spectrum antibiotic.

*film-coated tablets*

16 x 500 mg

16 x 1000 mg

*powder for oral suspension*

100 ml (250 mg / 5 ml)

## ● AZITROMICIN BELUPO

## AZITHROMYCIN

Semi-synthetic macrolide antibiotic.

*film-coated tablets*

6 x 250 mg

3 x 500 mg

● BELOXIM<sup>®</sup>

## CEFUROXIME

Antibacterial for systemic use in adults and children, second-generation cephalosporin.

*film-coated tablets*

10 x 500 mg

## ● CEFALEKSIN BELUPO

## CEFALEXIN

Semi-synthetic cephalosporin bactericidal antibiotic.

*capsules*

16 x 500 mg

*powder for oral suspension*

100 ml (250 mg / 5 ml)

● CIFLOX<sup>®</sup>

## CIPROFLOXACIN

Synthetic antibiotic belong to a group of fluoroquinolones. Broad-spectrum antibiotic that is active against both Gram-positive and Gram-negative bacteria.

*film-coated tablets*

10 x 250 mg

10 x 500 mg

*solution for infusion*

5 x 100 ml (2 mg / 1 ml)

## ● DOKSICIKLIN BELUPO

## DOXYCYCLINE HYCLATE

Tetracycline wide-spectrum antibiotic.

*capsules*

6 x 100 mg

25 x 100 mg

## ERITROMICIN BELUPO

ERYTHROMYCIN ESTOLATE

Macrolide wide-spectrum antibiotic.

*capsules*

16 x 250 mg

## GENTAMICIN BELUPO

GENTAMICIN SULPHATE

Aminoglycoside wide-spectrum antibiotic, for intramuscular and intravenous use.

*solution for injections*

10 x 2 ml (80 mg / 2 ml)

10 x 2 ml (120 mg / 2 ml)

## HERPLEXIM<sup>®</sup>

ACYCLOVIR

Highly selective antiviral for treatment of infections caused by herpes simplex and varicella-zoster viruses.

*film-coated tablets*

35 x 400 mg

## ITRAC<sup>®</sup> 3

ITRACONAZOLE

Broad-spectrum oral antimycotic.

*capsules*

4 x 100 mg

28 x 100 mg

## KLAVOBEL<sup>®</sup> BID

AMOXICILIN TRIHYDRATE + POTASSIUM CLAVULANATE

Semi-synthetic penicillin antibiotic it is used to treat a wide range of infections caused by bacteria.

*film-coated tablets*

14 x (875 + 125) mg

## MAKCIN<sup>®</sup>

CLARITHROMYCIN

Broad-spectrum macrolide antibiotic.

*film-coated tablets*

14 x 500 mg

## MAKCIN<sup>®</sup> SR

CLARITHROMYCIN

Broad-spectrum macrolide antibiotic.

*prolonged-release tablets*

7 x 500 mg

## MEDAZOL<sup>®</sup>

### METRONIDAZOLE

Antiinfective, antiprotozoal used for treatment of anaerobic infections, protozoal diseases and for prophylaxis in surgical and gynecological procedures.

*solution for infusion*

100 ml (0.5%)

## NINUR<sup>®</sup>

### NITROFURANTOIN

An urinary anti-infective. It is used to treat specific urinary tract infections.

*capsules*

30 x 50 mg

## NOFLOX<sup>®</sup>

### NORFLOXACIN

Broad-spectrum, quinolone bactericidal agent indicated for the treatment of upper and lower, complicated and uncomplicated, acute and chronic urinary tract infections.

*film-coated tablets*

20 x 400 mg

## SILAPEN<sup>®</sup>K

### PHENOXYMETHYLPENICILLIN POTASSIUM

Semi-synthetic penicillin antibiotic, particularly suitable for treatment of streptococcal infections.

Each film-coated tablet contains 1,000,000 IU corresponding to 654 mg of Phenoxymethylpenicillin Potassium or 590 mg Phenoxymethylpenicillin.

Each film-coated tablet contains 1,500,000 IU corresponding to 981 mg of Phenoxymethylpenicillin Potassium or 885 mg Phenoxymethylpenicillin.

Each 5 ml of oral solution contains 250 mg of Phenoxymethylpenicillin as Phenoxymethylpenicillin Potassium Ph. Eur.

*film-coated tablets*

30 x 1,000,000 IU

30 x 1,500,000 IU

*powder for oral solution*

100 ml (250 mg / 5 ml)

## SULOTRIM<sup>®</sup>

### SULFAMETHOXAZOLE AND TRIMETHOPRIM

Combined antibacterial agent having two active components with a synergetic effect.

*tablets*

20 x (100 + 20 mg)

20 x (400 + 80 mg)

*oral suspension*

100 ml (200 + 40 mg / 5 ml)

## SULOTRIM<sup>®</sup> FORTE

### SULFAMETHOXAZOLE AND TRIMETHOPRIM

Combined antibacterial agent having two active components with a synergetic effect.

*tablets*

20 x (800 + 160 mg)

## ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS

● ASTREA<sup>®</sup>

## IMATINIB

Inhibits is a protein-tyrosine kinase inhibitor that potently inhibits the activity of the Bcr-Abl tyrosine kinase, as well as several tyrosine kinase receptors. Shows antitumor activity in vivo.

*film-coated tablets*

60 x 100 mg

30 x 400 mg

● BIXALAN<sup>®</sup>

## BICALUTAMID

Antineoplastic agent, antiandrogen, non-steroidal androgen receptor inhibitor. It competitively inhibits the action of androgens by binding to cytosol androgen receptors in the target tissue.

*film-coated tablets*

28 x 50 mg

28 x 150 mg

## ● DOCETAKSEL BELUPO

## DOCETAXEL

Antineoplastic agent, taxan, used as a single agent or in combination with chemotherapeutic drugs as indicated depending on specific cancer type and stage, such as breast cancer, non-small lung cancer, prostate cancer, gastric adenocarcinoma, head and neck cancer.

*concentrate for solution for infusion*

20 mg / 0.5 ml

80 mg / 2 ml

● LETRILAN<sup>®</sup>

## LETROZOLE

Nonsteroidal competitive inhibitor of the aromatase enzyme system; it inhibits the conversion of androgens to estrogens. Letrozole inhibits the aromatase enzyme by competitively binding to the heme of the cytochrome P450 subunit of the enzyme, resulting in a reduction of estrogen biosynthesis in all tissues.

*film-coated tablets*

30 x 2.5 mg

## ● STRAZOLAN®

### ANASTROZOLE

Potent and highly selective nonsteroidal aromatase inhibitor. Anastrozole selectively inhibits the conversion of androgens to estrogens by inhibiting the aromatase enzyme complex through competitive binding to the haeme of the cytochrome P450 (CYP450) unit of the enzyme. Suppression of estrogen biosynthesis in all tissues reduces serum concentrations of circulating estrogens, including estrone, estradiol, and estrone sulphate with a subsequent beneficial effect in breast cancer.

*film-coated tablets*

28 x 1 mg

## ● TAKROLIMUS BELUPO

### TACROLIMUS

Highly potent immunosuppressive agent. Indicated in prophylaxis of transplant rejection in liver, kidney or heart allograft recipients and treatment of allograft rejection resistant to treatment with other immunosuppressive medicinal products.

*hard capsules*

30 x 0.5 mg

60 x 1 mg

M

## MUSCULO-SKELETAL SYSTEM

## ● ALEDOX®

### ALENDRONATE SODIUM

Potent bisphosphonate for the prevention and the treatment of postmenopausal osteoporosis.

*tablets*

4 x 70 mg

12 x 70 mg

## ● ALOPURINOL BELUPO

### ALLOPURINOL

An antigout agent. The primary use of allopurinol is to treat hyperuricemia and its complications. Allopurinol does not alleviate acute attacks of gout, but is useful in chronic gout to prevent future attacks.

*tablets*

100 x 100 mg

30 x 200 mg

100 x 200 mg

## BONNA®

### RISEDRONATE SODIUM

Risedronate is a potent bisphosphonate in the treatment of established postmenopausal osteoporosis in the women, osteoporosis in the man and osteoporosis caused by medicines (glucocorticoides).

*film-coated tablets*

4 x 35 mg                      2 x 75 mg  
6 x 75 mg

## IBUPROFEN BELUPO

### IBUPROFEN

Non-steroidal anti-inflammatory agent (NSAID) with analgesic and antipyretic effect. Rapidly and efficiently removes severe, moderate and sudden pain in acute rheumatic conditions, in soft tissue injuries, acute disorders of the lumbar spinal column and after surgery.

*film-coated tablets*

30 x 400 mg  
30 x 600 mg  
30 x 800 mg

*oral suspension*

100 ml (100 mg / 5 ml)

## INDOMETACIN BELUPO

### INDOMETACIN

Non-steroidal anti-inflammatory agent (NSAID). Helps to decrease swelling and pain.

*hard capsules*

30 x 25 mg

## KNAVON®

### KETOPROFEN

Non-steroidal anti-inflammatory agent (NSAID). Used in inflammatory and degenerative rheumatic diseases to reduce joint swelling, to reduce pain both at rest and in motion, to shorten the duration of morning stiffness, enable greater flexibility and functional capacity. Also suitable for removing abdominal pain.

*capsules*

25 x 50 mg

*suppositories*

12 x 100 mg

## KNAVON® FORTE

### KETOPROFEN

Non-steroidal anti-inflammatory agent (NSAID). Used in inflammatory and degenerative rheumatic diseases to reduce joint swelling, to reduce pain both at rest and in motion, to shorten the duration of morning stiffness, enable greater flexibility and functional capacity. Also suitable for removing abdominal pain.

*film-coated tablets*

20 x 100 mg

## KNAVON<sup>®</sup> RETARD

### KETOPROFEN

Non-steroidal anti-inflammatory agent (NSAID). Used in inflammatory and degenerative rheumatic diseases to reduce joint swelling, to reduce pain both at rest and in motion, to shorten the duration of morning stiffness, enable greater flexibility and functional capacity. Also suitable for removing abdominal pain.

*prolonged-release tablets*

20 x 150 mg

## LUBOR<sup>®</sup>

### PIROXICAM

Non-steroidal anti-inflammatory agent (NSAID), which effectively removes edema, erythema, and tissue proliferation in the affected joint. Used as analgesic in supporting therapy.

*capsules*

20 x 10 mg

20 x 20 mg

*suppositories*

10 x 20 mg

## LUBORETA<sup>®</sup>

### PIROXICAM

A topical NSAID that relieves pain due to sprains, contusions, strained muscles and for treatment of particular rheumatism.

*cream*

50 g (10 mg / g)

## NEOFEN<sup>®</sup> 200 mg tablets

### IBUPROFEN

Non-steroidal anti-inflammatory agent (NSAID) with analgesic and antipyretic effect. Removes mild and moderate pain of various origin (headache, toothache, premenstrual and menstrual pain, pain caused by injury, rheumatic diseases, cold and influenza). Particularly suitable for treatment of persistent high temperature (over 39°C).

*film-coated tablets*

20 x 200 mg

30 x 200 mg

## NEOFEN<sup>®</sup> direkt

### IBUPROFEN

Non-steroidal anti-inflammatory agent (NSAID) with analgesic and antipyretic effect. Removes mild and moderate pain of various origin (headache, toothache, premenstrual and menstrual pain, pain caused by injury, rheumatic diseases, cold and influenza). Particularly suitable for treatment of persistent high temperature (over 39°C).

*orodispersible tablets*

10 x 200 mg

## NEOFEN<sup>®</sup> forte

IBUPROFEN

Non-steroidal anti-inflammatory agent (NSAID) with analgesic and antipyretic effect. Removes mild and moderate pain of various origin (headache, toothache, premenstrual and menstrual pain, pain caused by injury, rheumatic diseases, cold and influenza). Particularly suitable for treatment of persistent high temperature (over 39°C).

*film-coated tablets*

10 x 400 mg

20 x 400 mg

## NEOFEN<sup>®</sup> plus gel

IBUPROFEN

Non-steroidal anti-inflammatory agent (NSAID) for topical treatment. Relives pain due to sprains, contusions, strained muscles and for treatment of particular rheumatism and sports injuries.

*gel*

50 g (50 mg / g)      100 g (50 mg / g)

## NEOFEN<sup>®</sup> spray

IBUPROFEN

Non-steroidal anti-inflammatory agent (NSAID) for topical treatment. Relives pain due to sprains, contusions, strained muscles and for treatment of particular rheumatism and sports injuries.

*spray*

50 ml (50 mg / g)

## NEOFEN<sup>®</sup> syrup

IBUPROFEN

Non-steroidal anti-inflammatory agent (NSAID). It is used for treatment of mild to moderate pain, post-immunization pyrexia, rheumatic or muscular pain, headache, reduction of fever, sore throat, teething pain, toothache, minor aches and pains, symptoms of cold and influenza.

*syrup*

100 ml (100 mg / 5 ml)

## NEOFEN<sup>®</sup> 60 g suppositories

IBUPROFEN

Non-steroidal anti-inflammatory agent (NSAID). It is used for treatment of mild to moderate pain, post-immunization pyrexia, rheumatic or muscular pain, headache, reduction of fever, sore throat, teething pain, toothache, minor aches and pains, symptoms of cold and influenza in children age 3 months to 2 years.

*suppositories*

10 x 60 mg

## NEOFEN<sup>®</sup> 125 g suppositories

IBUPROFEN

Non-steroidal anti-inflammatory agent (NSAID). It is used for treatment of mild to moderate pain, post-immunization pyrexia, rheumatic or muscular pain, headache, reduction of fever, sore throat, teething pain, toothache, minor aches and pains, symptoms of cold and influenza in children age 2-6 years.

*suppositories*

10 x 125 mg



# N

## NERVOUS SYSTEM

### ● ARVIND®

#### LAMOTRIGINE

Anticonvulsant. Used alone or in combination with other medications in the treatment of seizures in adults and children with epilepsy and Lennox-Gastaut syndrome. Lamotrigine is also used in the treatment of bipolar disorder.

*tablets*

30 x 25 mg      30 x 50 mg      30 x 100 mg

### ● AZOLAR®

#### ARIPIIPRAZOLE

Antipsychotic indicated for the treatment schizophrenia and manic episodes in bipolar I. disorder.

*tablets*

30 x 10 mg      30 x 15 mg      30 x 30 mg

### ● CALIXTA®

#### MIRTAZAPINE

Tetracyclic antidepressant. Noradrenergic and specific serotonergic antidepressant (NaSSA).

*film-coated tablets*

30 x 15 mg                      30 x 30 mg                      30 x 45 mg

### ● CERSON®

#### NITRAZEPAM

Benzodiazepine hypnotic, indicated for short-term treatment of insomnia.

*tablets*

10 x 5 mg                      30 x 5 mg

### ● CITRAM®

#### ESCITALOPRAM

An antidepressant of the selective serotonin reuptake inhibitor (SSRI) class, noted for its high selectivity of serotonin reuptake inhibition. It is the S-stereoisomer (enantiomer) of the earlier drug citalopram. It is indicated for the treatment in adults with major depressive disorder, generalized anxiety disorder and social anxiety disorder.

*film-coated tablets*

28 x 10 mg

## EMINENS®

### ROPINIROLE

A dopaminergic antiparkinsonism agent. Ropinirole may be used alone (without levodopa) in the treatment of idiopathic Parkinson's disease. Addition of ropinirole to levodopa may be used to control "on-off" fluctuations and permit a reduction in the total daily dose of levodopa.

#### *film-coated tablets*

210 x 0.25 mg

21 x 1 mg

## EMINENS® SR

### ROPINIROLE

Non-ergoline D2/D3 dopamine agonist that alleviates dopamine deficiency by stimulating striatal dopamine receptors.

#### *prolonged-release tablets*

28 x 2 mg

28 x 4 mg

28 x 8 mg

## HALEA®

### SERTRALINE

An antidepressant. Selective serotonin reuptake inhibitor (SSRI) used to treat depression, obsessive-compulsive disorder, panic disorder, posttraumatic stress disorder (PTSD).

#### *film-coated tablets*

30 x 50 mg

## KATENA®

### GABAPENTIN

A gamma-aminobutyric acid analog. Gabapentin is indicated as therapy in the treatment of partial seizures in adults with epilepsy with and without secondary generalization.

#### *capsules*

20 x 100 mg

50 x 300 mg

50 x 400 mg

## LEVETIRACETAM BELUPO

### LEVETIRACETAM

Antiepileptic, indicated as monotherapy for adults and adolescents from 16 years of age or as add-on therapy for adults ( $\geq 18$  years) and adolescents (12 to 17 years).

#### *film-coated tablets*

60 x 250 mg

60 x 500 mg

60 x 1000 mg

## LIDOKAIN BELUPO

### LIDOCAINE

A local anesthetic and antiarrhythmic

#### *solution for injections*

5 x 5 ml (2%)

100 x 2 ml (2%)

#### *spray*

50 ml 10% (100 mg / ml)

50 ml 10% (100 mg / ml)

## LIDOKAIN-ADRENALIN BELUPO

LIDOCAINE, ADRENALIN

A local anesthetic and antiarrhythmic

***solution for injections***

100 x 2 ml (40 + 0.025 mg / 2 ml)

## LORSILAN<sup>®</sup>

LORAZEPAM

Benzodiazepine anxiolytic. Due to simple inactivation (no intermediate products) it is suitable for treatment of patients with liver disease and older patients.

***tablets***

20 x 2.5 mg                      30 x 1 mg

## LUMIDOL<sup>®</sup>

TRAMADOL

Opioidic analgesic for treatment of moderate to severe acute and chronic pain.

***hard capsules***

20 x 50 mg

***oral drops***

10 ml (100 mg / ml)

***solution for injections***

5 x 1 ml (50 mg / ml)

5 x 2 ml (100 mg / 2 ml)

## LUMIDOL<sup>®</sup> RETARD

TRAMADOL

Opioidic analgesic for treatment of moderate to severe acute and chronic pain.

***prolonged-release tablets***

30 x 100 mg                      30 x 200 mg

## LUPOCET<sup>®</sup> 500 mg tablets

ACETAMINOPHEN (PARACETAMOL)

Paracetamol is an antipyretic and analgesic. It is recommended to relieve pain and to reduce fever. It is particularly suitable for the treatment of patients sensitive to salicylates.

***tablets***

20 x 500 mg

## LUPOCET<sup>®</sup> 500 mg film-coated tablets

ACETAMINOPHEN (PARACETAMOL)

Paracetamol is an antipyretic and analgesic. It is recommended to relieve pain and to reduce fever. It is particularly suitable for the treatment of patients sensitive to salicylates.

***film-coated tablets***

18 x 500 mg

## LUPOCET<sup>®</sup> 1000 mg effervescent tablets

ACETAMINOPHEN (PARACETAMOL)

Lupocet effervescent tablets are intended for lowering of fever and relieving of pain of various origins. It is particularly suitable for the treatment of patients sensitive to salicylates.

***effervescent tablets***

8 x 1000 mg

## ● LUPOCET® forte 1000 mg film-coated tablets

ACETAMINOPHEN (PARACETAMOL)

Paracetamol is an antipyretic and analgesic. It is recommended to relieve pain and to reduce fever. It is particularly suitable for the treatment of patients sensitive to salicylates.

*film-coated tablets*

10 x 1000 mg

## ● LUPOCET® baby 120 mg suppositories

ACETAMINOPHEN (PARACETAMOL)

LUPOCET suppositories are indicated for children for reducing fever and relieving pain caused by dentition, toothache, headache, pain after surgery or injuries, muscle and joint pain.

The use of paracetamol in suppository form is particularly suitable for patients who cannot take the oral form (e.g. after surgery or due to nausea or vomiting).

*suppositories*

10 x 120 mg

## ● LUPOCET® 250 mg suppositories

ACETAMINOPHEN (PARACETAMOL)

LUPOCET suppositories are indicated for children for reducing fever and relieving pain caused by dentition, toothache, headache, pain after surgery or injuries, muscle and joint pain.

The use of paracetamol in suppository form is particularly suitable for patients who cannot take the oral form (e.g. after surgery or due to nausea or vomiting).

*suppositories*

10 x 250 mg

## ● LUPOCET® flu effervescent tablets

ACETAMINOPHEN (PARACETAMOL), CHLORPHENIRAMINE MALEATE, VITAMINE C

LUPOCET FLU relieves the symptoms of cold and flu such as running and itchy nose, running eyes, hoarseness, cough, headache, fever and fatigue.

1 effervescent tablet contains 300 mg acetaminophen (paracetamol), 1.41 mg chlorpheniramine and 250 mg ascorbic acid.

*effervescent tablets*

10 x (300 mg + 1.41 mg + 250 mg)

## ● LUPOCET® flu forte effervescent tablets

ACETAMINOPHEN (PARACETAMOL), CHLORPHENIRAMINE MALEATE, VITAMINE C

LUPOCET FLU forte relieves the symptoms of cold and flu such as running and itchy nose, running eyes, hoarseness, cough, headache, fever and fatigue.

1 effervescent tablet contains 500 mg acetaminophen (paracetamol), 1.41 mg chlorpheniramine and 250 mg ascorbic acid.

*effervescent tablets*

10 x (500 mg + 1.41 mg + 250 mg)

## ● LUPOCET<sup>®</sup> teen capsules

### ACETAMINOPHEN (PARACETAMOL)

Paracetamol is an antipyretic and analgesic. Its use is recommended to relieve pain and to reduce fever in the common cold or flu, and for other pains such as headache (including migraine and tension headache), toothache, neuralgia, back pain, rheumatic and muscle pain, dysmenorrhoea and sore throat. It is also recommended for symptomatic relief of pain in mild arthritis. It is particularly suitable for the treatment of patients sensitive to salicylates.

#### *capsules*

20 x 300 mg

## ● LUPOCET<sup>®</sup> junior syrup

### ACETAMINOPHEN (PARACETAMOL)

Paracetamol is indicated for children for lowering fever associated with common cold, flu, infectious childhood diseases (e.g. chicken pox, whooping cough, measles, mumps) or reactions to vaccinations, and for relieving toothache, headache, sore throat, dentition pain, pain after surgery or injuries, muscle and joint pain.

5 ml syrup contains 120 mg acetaminophen (paracetamol).

#### *syrup*

100 ml

## ● MISAR<sup>®</sup>

### ALPRAZOLAM

Benzodiazepine anxiolytic used to relieve anxiety, nervousness, and tension associated with anxiety disorders. Also used to treat panic disorders.

#### *tablets*

30 x 0.25 mg

30 x 0.5 mg

30 x 1 mg

## ● MISAR<sup>®</sup> SR

### ALPRAZOLAM

Benzodiazepine anxiolytic (see previous).

#### *prolonged-release tablets*

30 x 0.5 mg

30 x 1 mg

## ● NORMABEL<sup>®</sup>

### DIAZEPAM

Benzodiazepine anxiolytic, sedative and central miorelaxant. Used to relieve anxiety, nervousness, and tension associated with anxiety disorders. It is also used to treat certain types of seizure disorders and muscle spasms.

#### *film-coated tablets*

30 x 2 mg

30 x 5 mg

#### *tablets*

30 x 10 mg

#### *solution for injections*

10 x 2 ml (10 mg / 2 ml)

## ● OKSAZEPAM BELUPO

### OXAZEPAM

Benzodiazepine anxiolytic

#### *tablets*

30 x 10 mg

30 x 15 mg

30 x 30 mg

## PROMAZIN<sup>®</sup>

### PROMAZINE

Phenothiazine class of antipsychotics. Used as an adjunct to short-term management of moderate to severe psychomotor agitation. It is also used to treat agitation and restlessness in the elderly.

#### *coated tablets*

50 x 25 mg

50 x 100 mg

## PROSPERA<sup>®</sup>

### RISPERIDONE

Antipsychotic for the treatment of schizophrenia and mania associated with bipolar disorder.

#### *film-coated tablets*

20 x 1 mg

60 x 2 mg

60 x 3 mg

60 x 4 mg

## Q-PIN<sup>®</sup>

### QUETIAPINE

Antipsychotic for the treatment of schizophrenia and mania associated with bipolar disorder.

#### *film-coated tablets*

60 x 25 mg

60 x 100 mg

60 x 200 mg

## SULPIRID BELUPO

### SULPIRIDE

Antipsychotic; antidepressant, anxiolytic.

#### *capsules*

30 x 50 mg

12 x 200 mg

## TIRAMAT<sup>®</sup>

### TOPIRAMATE

A carbonic anhydrase inhibitor anticonvulsant. It is used to treat seizures in adults or children with epilepsy. Topiramate is indicated in adults for the prophylaxis of migraine headache.

#### *film-coated tablets*

60 x 25 mg

60 x 50 mg

60 x 100 mg

## URUTAL<sup>®</sup>

### BETAHISTINE

Antivertigo-hemokinetic and otolitic.

#### *tablets*

100 x 8 mg

60 x 16 mg

20 x 24 mg

50 x 24 mg

## VAIRA<sup>®</sup>

### OLANZAPINE

Antipsychotic used to treat the symptoms of psychotic conditions such as schizophrenia and bipolar disorder.

*film-coated tablets*

28 x 5 mg

28 x 10 mg

## VAIRA<sup>®</sup>-V

### OLANZAPINE

Antipsychotic used to treat the symptoms of psychotic conditions such as schizophrenia and bipolar disorder.

*orodispersible tablets*

28 x 5 mg

28 x 10 mg

## ZARACET<sup>®</sup>

### TRAMADOL / ACETAMINOPHEN (PARACETAMOL)

Combination of opioidic analgesic and analgoantipiretic for symptomatic treatment of moderate and severe pain.

*film-coated tablets*

10 x (37.5 mg + 325 mg)

20 x (37.5 mg + 325 mg)

30 x (37.5 mg + 325 mg)

100 x (37.5 mg + 325 mg)

## P

## ANTIPARASITIC PRODUCTS

● MEDAZOL<sup>®</sup>

## METRONIDAZOLE

Antiinfective, antiprotozoal used for treatment of anaerobic infections, protozoal diseases and for prophylaxis in surgical and gynecological procedures.

*tablets*

20 x 250 mg

10 x 400 mg

20 x 400 mg

## R

## RESPIRATORY SYSTEM

● ALERDIN<sup>®</sup>

## DESLORATADINE

Desloratadine is non-sedating long-acting histamine antagonist with selective peripheral H<sub>1</sub>-receptor antagonist activity.

*orodispersible tablets*

10 x 5 mg

30 x 5 mg

*oral solution*

60 ml (0.5 mg / ml)

120 ml (0.5 mg / ml)

● BELODIN<sup>®</sup> / BELODIN<sup>®</sup>A

## LORATADINE

Antihistaminic for systematic use.

Loratadine is used in the treatment of allergy symptoms, such as:

- allergic rhinitis (seasonal and perennial)
- chronic urticaria and other allergic skin disturbances, including neurodermatitis.

*tablets (BELODIN)*

30 x 10 mg

*tablets (BELODIN A)*

7 x 10 mg

14 x 10 mg



## ● MONLAST<sup>®</sup>

### MONTELUKAST

A leukotriene receptor antagonist (LTRA) used for the maintenance treatment of asthma and to relieve symptoms of seasonal allergies. Montelukast blocks the action of leukotriene D4 on the cysteinyl leukotriene receptor CysLT1 in the lungs and bronchial tubes. This reduces the bronchoconstriction otherwise caused by the leukotriene, and results in less inflammation. It is not useful for the treatment of acute asthma attacks.

*chewable tablets*  
28 x 5 mg

*film-coated tablets*  
28 x 10 mg

## ● RINIL<sup>®</sup> 0.5 mg / ml

### XYLOMETAZOLINE

Application to the nasal passages for children at the age of 2 – 10 years. For the symptomatic relief of nasal congestion, perennial and allergic rhinitis, sinusitis.

*nasal spray, solution*  
10 ml

## ● RINIL<sup>®</sup> 1 mg / ml

### XYLOMETAZOLINE

Application to the nasal passages for adults and children at the age  $\geq 10$  years. For the symptomatic relief of nasal congestion, perennial and allergic rhinitis, sinusitis.

*nasal spray, solution*  
10 ml

## FOOD FOR SPECIAL MEDICAL PURPOSES

## ● NUTRIXA<sup>®</sup>

Only for use under medical supervision

Complete and balanced nutrition. For the dietary management of patients with malnutrition, or at risk of developing malnutrition. High calorie, high protein liquid food with added fiber for a complete, balanced diet. For oral use.

200 ml of high protein liquid food corresponds to 250 kcal.

1 ml of high protein liquid food corresponds to 1.25 kcal.

200 ml of high protein liquid food contain 16 g protein.

Flavour vanilla / Flavour chocolate

200 ml

# HERBAL MEDICINES, FOOD SUPPLEMENTS AND COSMETICS



## HERBAL MEDICINES

### AKTIVIN® - H

*Hyperici herbae extractum siccum*

AKTIVIN H is herbal antidepressant used to induce good mood in mild depression and symptoms such as restlessness, anxiety and irritability.

1 capsule contains: 225 mg extract (as dry extract) of *Hypericum perforatum* L., herba (dried flowering tops of St. John's wort) corresponding to 0.5 mg of total hypericin

*30 capsules, 60 capsules*

### GINKGO

*Ginkgo bilobae folii extractum siccum*

GINKGO film-coated tablets are used to relieve problems associated with venous and capillary blood flow in the following cases:

- brain circulation disorder manifested by gradual deterioration of brain functions and the following symptoms: vertigo, tinnitus, headache, diminishing memory, insomnia, bad mood and fear.
- restriction of circulation in cervical vertebrae leading to diminishing hearing ability;
- circulatory insufficiency in the extremities manifested by: loss of sensation, pins and needles, cramps, numbness, coldness or burning etc.
- varicose veins and hemorrhoids.

1 film-coated tablet contains: 40 mg extract (as dry extract) of *Ginkgo biloba* L., folium (ginkgo leaf) (40-50:1) standardized to at least 8.8 – 10.8 mg of flavonoids expressed as flavonoid glycosides, 1.12 – 1.36 mg of ginkgolid A,B,C and 1.04 – 1.28 mg of bilobalid.

*60 film-coated tablets x 40 mg*

*90 film-coated tablets x 40 mg*

## ● HEDERAN<sup>®</sup> syrup

*Hederae helicis folii extractum siccum*

Hederan syrup is used as an expectorant in adults and children at the age of 2 and over.

1 ml of syrup contains 3.675 mg of Ivy dry extract (*Hedera helix* L., folium).

*125 ml syrup*

*200 ml syrup*

## ● HEDERAN<sup>®</sup> film-coated tablets

*Hederae helicis folii extractum siccum*

Hederan tablets are used as an expectorant in adults, adolescents and children aged 5 years and over.

1 film coated tablet contains: 35 mg of Ivy dry extract (*Hedera helix* L., folium).

*20 film-coated tablets*

## ● SILYMARIN

*Silybi mariani fructi extractum siccum*

SILYMARIN capsules are used to alleviate liver dysfunction.

1 capsule contains 156.7-191.6 mg of refined and standardised dry extract of the milk thistle fruit, *Silybum marianum* (L.) Gaertn which corresponds to 100 mg of silymarin expressed as silibinin.

*30, 60 capsules*

## ● SILYMARIN FORTE

*Silybi mariani fructi extractum siccum*

SILYMARIN film-coated tablets are used to alleviate liver dysfunction due to chronic hepatitis B and C, alcoholic liver disease and cirrhosis.

1 capsule contains 327.59 – 362.07 mg of refined and standardised dry extract of the milk thistle fruit, *Silybum marianum* (L.) Gaertn which corresponds to 200 mg of silymarin expressed as silibinin.

*30 film-coated tablets*

# FOOD SUPPLEMENTS

## **BELMIRAN<sup>®</sup> DAN (BELMIRAN<sup>®</sup> DAY)**

*Passiflora incarnata*

BELMIRAN DAN tablets contain passion flower, which contributes to relaxation of the body.

1 film-coated tablet contains: 100 mg of passion flower dry extract (*Passiflorae incarnatae*).

*20 film-coated tablets*

## **BELMIRAN<sup>®</sup> SAN (BELMIRAN<sup>®</sup> NIGHT)**

*Valeriana officinalis, Humulus lupulus, Passiflora incarnata*

BELMIRAN SAN tablets help shorten the time needed to initiate and maintain sleep.

1 film-coated tablet contains: 125 mg valerian root dry extract (*Valeriana officinalis*), 30 mg hops strobile dry extract (*Humulus lupulus*) and 80 mg passion flower dry extract (*Passiflora incarnata*).

*20 film-coated tablets*

## **BEVITAL<sup>®</sup> SUN EXTRA**

*Vitamin C, Vitamin E, Lycopene,  $\beta$ -carotene, Lutein*

BEVITAL SUN EXTRA capsules contain vitamin C, which contributes to normal collagen formation for the normal function of the skin. Contain vitamin C and vitamin E which contributes to protection of cells against oxidative stress.

1 capsule contains: 100.0 mg of vitamin C (125% RDA), 30.0 mg of vitamin E (250% RDA), 8.0 mg of lycopene, 5.67 mg of  $\beta$ -carotene, 0.2 mg of lutein.

*30 capsules*

## **FERSAN<sup>®</sup> FORTE tablets**

*Carobonyl iron, Vitamin C, Folic acid*

FERSAN FORTE tablets contain iron, vitamin C and folic acid. Iron contributes to normal red blood cell and hemoglobin formation and normal cognitive development of children. Folic acid contributes to normal blood formation and cell division. Vitamin C increases iron absorption.

1 tablet contains: 30 mg (214% RDA) carbonyl iron, 180 mg (225% RDA) vitamin C and 600  $\mu$ g (300% RDA) folic acid.

*30 tablets*

## FERSAN<sup>®</sup> JUNIOR

*Carobonyl iron, Vitamin C, Folic acid*

FERSAN JUNIOR liquid is dietary supplement containing iron which contributes to normal: function of the immune system, cognitive function, metabolism, red blood cells and hemoglobin, transport of oxygen in the body. It contributes in reducing fatigue and exhaustion and has a role in the process of cell division.

30 ml of liquid supplement contains 814.7 mg of ferric pyrophosphate which corresponds to 210 mg elemental iron.

*drops (30 ml)*

## FAVORA<sup>®</sup> MARSHMALLOW

*Malva sylvestris, Althaea officinalis*

FAVORA MARSHMALLOW contributes to relieve irritation of the throat and pharynx.

125 ml of liquid food supplement contains: 2.6 g of mallow dry extract (*Malva sylvestris*) and 0.525 g of Marshmallow root dry extract (*Althaea officinalis*).

*125 ml food supplement*

## FAVORA<sup>®</sup> THYME

*Pimpinella anisum, Thymus vulgaris, vitamin C, Primula veris, Primula elatior*

FAVORA THYME contributes to relieve irritation of upper respiratory tract.

125 ml of liquid food supplement contains: 1.564 g slurry anis fruit extract (*Pimpinella anisum*), 0.317 g slurry thyme extract (*Thymus vulgaris*), at least 0.244 g of vitamin C (ascorbic acid) and 0.208 g slurry aqueous primula root extract (*Primula veris*, *Primula elatior*).

*125 ml food supplement*

## HedAKUT granules

*N-acetylcysteine, Vitamin C, Zinc, Marshmallow root dry extract (Althaea officinalis), blue eucalyptus (Eucalyptus globulus)*

HedAKUT granules contain marshmallow and blue eucalyptus that contribute to the relief in the case of irritation of the upper respiratory tract. HedAKUT granules contain vitamin C and zinc, which contribute to the normal function of the immune system.

1 sachet contains: 200 mg of N-acetylcysteine, 60 mg of Vitamin C (75% RDA), 50 mg of Marshmallow dry extract, 10 mg of Zinc (100% RDA), 3 mg of Eucalyptus dry extract

*36 g (12 sachets x 3 g)*

## MAXI OMEGA-3

*OMEGA-3 fatty acids*

MAXI OMEGA-3 capsules contain EPA / DHA which contribute to the normal function of the heart.

1 capsule contains 1000 mg of purified and standardized fish oil with a minimum of 30% of omega-3 fatty acids (EPA / DHA), including 18% of eicosapentaenoic acid (EPA) and 12% of docosahexaenoic acid (DHA).

*100 capsules*

## UROSAL<sup>®</sup>

*Sabalıs serrulatae fructi extractum siccum, Solidaginis virgaureae herbae extractum siccum, Urticae radıcıs extractum siccum*

UROSAL capsules relieve prostate problems, have an antiseptic effect on the urinary tract, and reduce irritation and pain related to urinating.

1 capsule contains 25 mg dry extract of the saw palmetto fruit, *Serenoa repens* (Bartram) Small. (*Sabal serrulata* (Michaux) Nichols); 15 mg dry extract of European goldenrod *Solidago virgaurea* L.; 10 mg dry extract of the root of stinging nettle *Urtica dioica* L. and/or dwarf nettle *Urtica urens* L.

*50 capsules*

## UROSAL<sup>®</sup> MAN

*Sabalıs serrulatae fructi extractum spissum*

UROSAL MAN capsules are intended for adult male persons.

1 capsule contains 240 mg dry extract *Sabal serrulata*.

*30 capsules*

## UROSAL<sup>®</sup> LADY

*Uvae ursi folii extractum siccum, Solidaginis virgaureae herbae extractum siccum, Urticae radıcıs extractum siccum.*

1 film-coated tablet contains: 400 mg dry extract of bearberry leaf (*Uvae ursi folii extractum siccum*), 15 mg dry extract of the European goldenrod herb (*Solidaginis virgaureae herbae extractum siccum*), 10 mg dry extract of stinging nettle (*Urticae radıcıs extractum siccum*).

*30 film-coated tablets*

## VITASENG<sup>®</sup>

*Panax ginseng radıcıs extractum*

VITASENG film-coated tablets contain ginseng root which contributes to normal erectile function and helps maintain vitality.

1 film-coated tablet contains 100 mg standardized dry extract of ginseng root, *Panax ginseng* (8% ginsenosides).

*30 film-coated tablets*

# COSMETICS

## EUSIN<sup>®</sup> oil

*Eusyn globulus oil, Mentha piperita oil, Melaleuca leucadendra oil, Menthol, Juniperus communis oil, Methyl salicylate, Eugenia caryophyllus oil*

EUSIN oil is used in baths.

EUSIN oil contains mixture of essential oils: eucalyptus oil (Eucalyptus globulus), peppermint oil (Mentha piperita), cajuput oil (Melaleuca leucadendra), juniper oil (Juniperus communis) and clove oil (Eugenia caryophyllus). It contains also: Methyl salicylate, Menthol, limonene and eugenol.

*drops (10 ml)*

## FLOCETA<sup>®</sup> gel

*Chamomilla recutita extract, Calendula officinalis extract, Aluminium-acetotartarate*

FLOCETA gel is used for skin care in the case of skin irritation due to sun exposure, shaving, waxing, excessive sweating and other skin irritations.

FLOCETA gel contains: aqua, propylene glycol, tetrahydroxypropyl ethylenediamine, aluminium acetate-tartrate solution, chamomilla recutita flower extract, calendula officinalis flower extract, carbomer, polysorbate 80, methylparaben, disodium EDTA, propylparaben.

*gel (20 g)*

## FLOCETA<sup>®</sup> lotion

*Chamomilla recutita extract, Calendula officinalis extract, Aluminium-acetotartarate, bisabolol*

FLOCETA lotion is used for skin care in the case of skin irritation due to sun exposure, shaving, waxing, excessive sweating and other skin irritations.

FLOCETA lotion contains: aqua, propylene glycol, aluminium acetate-tartrate solution, chamomilla recutita flower extract, calendula officinalis flower extract, polysorbate 80, methylparaben, propylparaben, sodium hydroxide, xanthan gum, bisabolol.

*lotion, spray for irritated skin (100 ml)*

## FLOCETA<sup>®</sup> soft gel

*Chamomilla recutita flower extract, Calendula officinalis flower extract, Hamamelis virginiana bark/leaf/twig extract*

FLOCETA SOFT gel soothes irritated, red and sensitive skin and gives a feeling of comfort, protects and regenerates the skin after exposure to external factors. Also convenient for the care of sensitive baby skin.

FLOCETA SOFT gel contains: aqua, chamomilla recutita flower extract, calendula officinalis flower extract, hamamelis virginiana bark/leaf/twig extract, glycerin, propylene glycol, tetrahydroxypropyl ethylenediamine, carbomer, ethylhexylglycerin, disodium EDTA, polysorbate 80.

*gel (15 g)*

 **BELOBAZA<sup>®</sup>****SOFT PARAFIN AND FAT PRODUCTS**

Ambiphilic base of carefully balanced ingredients for skin treatment, protection and care, and for preparation of magistral medicines (can be combined with almost any dermatotherapeutic).

*cream*

2 kg

4 kg